

angle of central fascia; some slight strigulation above tornus: cilia white, on termen with a grey subbasal line and outer third grey, on apex wholly dark grey. Hindwings thinly scaled, bluish-white; an apical blotch of grey suffusion; cilia whitish, suffused grey round apex.

PERU, Agualani, 9000 feet, June (*Ockenden*); 1 ex. (Coll. Paravicini).

*Tortrix apochranta*, n. sp.

♂. 16 mm. Head grey-whitish. Thorax grey-whitish somewhat sprinkled grey. Forewings suboblong, costa anteriorly moderately arched, edge slenderly reflexed on basal sixth, apex obtuse, termen slightly rounded, little oblique; pale whitish-ochreous-grey, with very faint reddish tinge, some thinly scattered grey strigulae; markings dark grey tinged reddish; central fascia moderate, rather oblique, marked blackish in disc, becoming nearly obsolete on dorsal third; costal patch moderately large, semioval: cilia pale grey, becoming whitish towards tornus. Hindwings pale greyish, strigulated grey posteriorly; cilia whitish.

BRAZIL, S. Leopoldo; 1 ex. (Coll. Joannis).

*Tortrix endopyrrha*, n. sp.

♂. 15–16 mm. Head, palpi grey. Antennal ciliations 1. Thorax fuscous. Forewings suboblong, costa anteriorly moderately arched, with subtriangular fold enlarged by rough projecting scales from near base to  $\frac{1}{4}$ , apex obtuse-pointed, termen slightly sinuate, little oblique; brown, bases of scales whitish: cilia pale greyish-ochreous. Hindwings dull fulvous-orange; cilia pale orange-ochreous.

UGANDA, Bulambuli, 9000 feet, July, August (*G. L. R. Hancock*); 4 ex. (type Brit. Mus.).

*Xenothictis atriflora*, n. sp.

♂. 19 mm. Head, palpi, thorax pale brownish-ochreous, a small brown spot on middle of tegulae. Antennal ciliations 1. Forewings moderate, posteriorly dilated, costa gently arched, apex obtuse, termen rounded, little oblique; pale ochreous, slightly speckled rufous; a blackish dot on base of costa, and two or three in disc near base; an irregular-triangular black blotch at about  $\frac{1}{4}$  from base below middle; upper half of central fascia black, rising from costa at  $\frac{2}{5}$ , very oblique, narrow and gradually dilated to middle of disc, where it is prolonged posteriorly into a rounded lobe; three or four black-brown spots or dots on costa posteriorly, and one near end of fascia: cilia pale ochreous. Hindwings grey, faintly darker-mottled; cilia pale grey.

Fiji, Vunidawa, December (*Phillips*); 1 ex. (Brit. Mus.). In rather imperfect condition, but very distinct.

*Eulia homosacta*, n. sp.

♂. 21 mm. Head white. Palpi  $2\frac{1}{4}$ , white, suffused pale brownish externally. Antennal ciliations  $1\frac{1}{2}$ . Thorax white, a dark grey spot on base of tegulae. Forewings elongate-triangular, costa slightly arched, apex obtuse-pointed, termen straight, rather oblique; white, coarse scattered light grey strigulae; markings fuscous, coarsely strigulated dark grey; an irregularly hemispherical blotch on base of costa reaching half across wing; a large elongate-triangular patch extending on costa from  $\frac{1}{4}$  to  $\frac{4}{5}$ , and reaching somewhat more than half across wing, beneath its apex in disc a small spot, and one beyond it on costa; a slender irregular terminal fascia: cilia whitish-grey, a grey subbasal shade, on costa white. Hindwings grey-whitish, obscurely mottled light grey on apical third and towards termen; cilia whitish, a grey subbasal shade.

ECUADOR, Hacienda Cayandeled, Prov. Rio Bamba, 4200 feet, February; 1 ex. (Coll. Paravicini). Near *sagmatica*, with very similar markings, but the different form of forewings (in *sagmatica* broader, with apex and termen rounded) forbids their identity.

*Paraptila hydrochoa*, n. sp.

♂. 17 mm. Head white, crown faintly pinkish, sides grey. Palpi greyish. Thorax white, erect lateral tufts mixed grey. Forewings moderate, posteriorly dilated, costa gently arched, apex obtuse, termen straight, somewhat oblique; white mixed light grey; basal patch light grey, edge formed by a suffused brownish line running from costa at  $\frac{1}{4}$  to middle of dorsum, slightly angular below middle; central fascia represented by a large greyish quadrilateral blotch edged fuscous extending on costa from before middle to  $\frac{3}{4}$  and reaching half across wing, in disc suffused fuscous posteriorly, its posterior angle extended as a rather curved streak to dorsum just before tornus; dorsal area between basal patch and central fascia irregularly clouded grey; some fine dark fuscous strigulation towards termen: cilia white. Hindwings grey-whitish, greyer and somewhat mottled grey posteriorly; a rather long slender ochreous-whitish subdorsal hair-pencil from base; cilia whitish.

ECUADOR, Hacienda Cayandeled, Prov. Rio Bamba, 4200 feet, February; 1 ex. (Coll. Paravicini). Although this has the three characteristic thoracic tufts of *Paraptila*, it appears superficially rather to belong to *Eulia*.

*Schœnotenes octosticta*, n. sp.

♂ 18 mm. Head, thorax white. Palpi white, second joint fuscous posteriorly except on apical third. Antennæ simple. Forewings suboblong, costa gently arched, apex obtuse, termen nearly straight, little oblique; ochreous-white, with faint more ochreous-tinged scattered strigæ, and some tufts of raised scales; eight small black dots on costa; two minute black raised dots on fold towards base, a small mark in disc before middle, and three or four minute dots in disc towards termen: cilia white. Hindwings and cilia white.

BRITISH NEW GUINEA, Owgarra (Meek); 1 ex. (Coll. Paravicini).

**LOPHOPRORA, n. g.**

Palpi very long, porrected, with dense rough projecting scales diminishing to apex, terminal joint short, nearly concealed. Antennæ simple. Thorax with double posterior crest. Forewings with tufts of scales; 2 from near  $\frac{2}{3}$ , 3 and 4 approximated from angle, 7 to costa. Hindwings without cubital pecten; cell  $\frac{1}{3}$ , 2-4 separate and approximated towards base, 5 parallel, 6 and 7 approximated at base.

Allied to *Spatalistis*.

**Lophoprora cyanostacta, n. sp.**

♂. 13 mm. Head, thorax fuscous. Palpi fuscous, suffused leaden-metallic above. Forewings moderate, slightly dilated, costa gently arched, rather bent and with projecting scales at  $\frac{3}{4}$ , apex pointed, termen concave, rather oblique; fuscous; a blotch of blue-leaden-metallic suffusion beneath costa at base, and some irregular spots forming antemedian, postmedian, and præterminal series; a large blackish tuft beneath fold at  $\frac{2}{5}$ , several smaller tufts scattered in disc, and two transversely placed above dorsum beyond middle; a pale yellowish costal streak marked ferruginous from  $\frac{1}{4}$  to  $\frac{3}{4}$ : cilia light yellow, base ferruginous round costa and apex, on costal prominence and tornus fuscous. Hindwings ochreous-yellow; a fuscous apical patch occupying about  $\frac{1}{3}$  of wing; cilia grey, round dorsum and tornus whitish-ochreous.

BRITISH NEW GUINEA, Kumusi R. (Meek); 1 ex. (Coll. Paravicini).

**Spatalistis droserantha, n. sp.**

♀. 15 mm. Head, thorax light yellow, mixed or mottled pale lilac-brownish. Palpi pale yellow, three oblique pale brownish bands. Forewings moderate, posteriorly dilated, costa gently arched, apex obtuse-pointed, termen sinuate, rather oblique; clear yellow, irregularly reticulated pale ochreous-brownish, with numerous raised pale violet-metallic dots in irregular transverse series; three or four larger blue-leaden dots beneath costa anteriorly; an incurved pale ochreous-brownish streak with several larger dots from middle of dorsum to costa about  $\frac{4}{5}$ : cilia whitish-yellow. Hindwings whitish-ochreous becoming whitish anteriorly; cilia ochreous-whitish.

ASSAM, Shillong, 5000 feet, May (Fletcher); 1 ex. Allied to *hormota*.

**Sparganothis elimata, n. sp.**

♂. 21 mm. Head, thorax ochreous-brown. Palpi  $2\frac{1}{2}$ , fuscous. Antennal ciliations  $1\frac{1}{2}$ . Forewings suboblong, costa anteriorly moderately, posteriorly slightly arched, apex obtuse, termen slightly rounded, hardly oblique; ochreous-brown, irregularly scattered indistinct small blackish dots: cilia ochreous-brown. Hindwings grey-whitish; cilia grey-whitish, a grey subbasal line.

ECUADOR, Hacienda Cayandeled, Rio Bamba, 4200 feet, February; 1 ex. (Coll. Paravicini).

*Argyrotoxa venatrix*, n. sp.

♂. 10 mm. Head grey. Palpi ochreous-whitish, terminal joint with dark fuscous ring. Thorax bluish-green. Forewings elongate-oblong, costa gently arched, apex obtuse, termen slightly rounded, rather oblique; bluish-green, terminal fourth light blue-grey, costal edge whitish-yellow marked red beneath, with several small blackish spots and dots; dorsum slenderly grey, edge partly whitish from  $\frac{1}{4}$  to  $\frac{3}{4}$ ; a scarlet mark on base of dorsum, a dot towards costa near base, an angulated line from near this to dorsum at  $\frac{1}{3}$ , a rather irregular slightly curved line from a black costal dot at  $\frac{1}{5}$  to near dorsum beyond middle, and a line from a small black costal spot before middle, right-angled in disc and interrupted above angle, to dorsum at  $\frac{3}{4}$ ; a fine whitish-ochreous terminal line, marked blackish round apex, terminating in an upwards-oblique pale ochreous whitish-edged lobe from just above tornus, marked with a black apical spot posteriorly: cilia pale leaden-grey mixed whitish. Hindwings dark fuscous; cilia grey.

CAMEROONS, Lolodorf (*Conradt*); 1 ex. (Coll. Paravicini). Much like *albardana* Snell (= *tricolor* Wals.), which however differs in neuration, and has a differently coloured thorax and other distinctions.

*Argyrotoxa erythrocyma*, n. sp.

♀. 15 mm. Head brown. Palpi brown, terminal joint grey-whitish with dark grey subapical ring. Thorax fuscous, tegulae blue-grey with a red streak. Forewings suboblong, costa moderately arched, apex obtuse, termen straight, hardly oblique, rounded beneath; light blue-grey, becoming blue on basal third; extreme costal edge whitish suffused reddish beneath and dotted blackish, with small blackish spots at middle and  $\frac{2}{3}$ ; base of dorsum red; six irregular somewhat interrupted rather curved oblique transverse red lines; a dark red somewhat interrupted marginal line round apex and upper part of termen: cilia whitish, base pale grey, on costa three blackish bars. Hindwings dark fuscous; cilia grey.

CAMEROONS, Lolodorf (*Conradt*); 1 ex. (Coll. Paravicini). Two other examples, also ♀ (1 from same locality, 1 from Johann-Albrechts Höhe Station) are probably varieties of the same species; these have groundcolour of forewings wholly greyish-violet, markings reddish-fuscous or partly deep ferruginous, similar but rather variable; in one of these a slender whitish line runs from near costa at  $\frac{1}{6}$  in an even curve through middle of disc to costa at  $\frac{5}{6}$ .

*Peronea ulmicola*, n. sp.

♀. 16 mm. Head brownish grey. (Palpi missing.) Thorax grey-brownish. Forewings suboblong, costa anteriorly strongly arched, slightly sinuate beyond middle, apex obtuse, termen distinctly sinuate, little oblique; grey mixed whitish; some blackish sprinkling in disc anteriorly; a tuft on fold at  $\frac{1}{4}$ ; a chestnut-brown triangular blotch extending on costa from before middle to  $\frac{4}{5}$  and reaching half across wing, its posterior edge confluent with a lighter brown

patch which extends to dorsum before tornus and covers posterior area except a narrow marginal fascia; some slight blackish sprinkling towards tornus: cilia fuscous sprinkled whitish. Hindwings grey, somewhat darker posteriorly; cilia light grey, a darker subbasal line.

MANCHURIA, bred October from larva rolling leaf of *Ulmus*; 1 ex. Easily distinguished from *boscana* by sinuate termen of forewings.

### *Peronea tephromorpha*, n. sp.

♂. 19 mm. Head, palpi dark grey. Thorax grey, slightly mixed brownish, with strong triangular crest. Forewings rather elongate-oblong, costa anteriorly moderately, posteriorly slightly arched, apex obtuse, termen rather obliquely rounded; ashy-grey, obscurely mixed light brownish, without apparent tufts: cilia grey, somewhat whitish-sprinkled. Hindwings grey; cilia pale grey.

MANCHURIA, bred from larva rolling leaf of *Ulmus*; 1 ex. Mr. August Busck, of the U.S. National Museum, who furnished me with this specimen, and has examined the genitalia of another example, states that they are very similar to those of *lipsiana*.

### *ANTIGRAPTIS*, n. g.

Palpi long, porrected, second joint expanded with scales towards apex above and beneath, terminal joint moderately long. Thorax without crest. Forewings with some very slight scale-tufts; 2 from  $\frac{2}{3}$ , 3 and 4 stalked from angle, 5 almost connate with stalk, 7 to termen, 8 and 9 stalked, connate with 7. Hindwings without cubital pecten; 3 and 4 connate, 5 curved, closely approximated at base, 6 and 7 closely approximated towards base.

Allied to *Peronea* and *Mictoneura*.

### *Antigraptis hemicrates*, n. sp.

♀. 22 mm. Head, thorax whitish-ochreous, shoulders fuscous. Palpi whitish-ochreous, second joint with indistinct fuscous band towards base, terminal joint with fuscous median ring. Forewings elongate-triangular, costa gently arched, apex obtuse-pointed, termen slightly sinuate, rather oblique; whitish-ochreous with faint pinkish tinge; blackish costal markings, viz. marks at and near base, a rather oblique fasciate streak at  $\frac{1}{5}$  reaching half across wing, transverse strigulae in disc before and beyond this, one or two costal marks at  $\frac{1}{4}$ , a truncate V-shaped blotch towards middle enclosing a costal dot, posterior arm slightly interrupted, two small spots towards  $\frac{3}{4}$ , a short transverse streak beyond this, and a dot just before apex; area below these markings irregularly marbled with brownish-ochreous, markings partly edged with fuscous strigulae or striæ; a fuscous terminal line: cilia light ochreous becoming whitish towards tips. Hindwings grey-whitish, with suffused light grey strigulae or striæ except towards base; cilia ochreous-whitish, a light greyish subbasal line.

DUTCH NEW GUINEA, Mt. Goliath, 5-7000 feet (*Meek*); 1 ex. (Coll. Paravicini). An interesting novel form.

*Cacoecia concavata*, n. sp.

♂. 29 mm. Head, palpi brown. Antennal ciliations  $2\frac{1}{2}$ . Thorax whitish-brownish, anterior margin dark brown. Forewings rather broad, costa anteriorly strongly arched, with moderate fold on basal  $\frac{1}{6}$ , posteriorly sinuate, apical area projecting obliquely upwards in a broad rounded prominence, termen slightly sinuate, somewhat oblique; brownish-ochreous, browner towards costa, faintly darker-strigulated, dorsum shortly strigulated dark grey; a ridge of raised ochreous-tinged scales beneath costa towards  $\frac{1}{3}$ : central fascia narrow, oblique, purplish-grey, ceasing below middle; costal patch semioval, purplish-grey, becoming dark fuscous towards costa inwardly, a terminal fascia of grey suffusion and darker strigulation becoming obsolete towards tornus: cilia ferruginous, becoming pale brownish-grey towards tornus. Hindwings whitish, some slight grey sprinkling towards termen; cilia whitish, on dorsum ochreous-tinged.

PERU, Agualani, Carabaya, 9000 feet, April (*Ockenden*), 1 ex. (Coll. Paravicini).

## ŒCOPHORIDÆ.

*Macrobathra neurocoma*, n. sp.

♂. 13-14 mm. Head, thorax dark bronzy-fuscous. Palpi very slender, dark fuscous, posteriorly white. Abdomen fuscous, dorsally suffused white towards base. Forewings elongate, costa gently arched, apex pointed, termen very obliquely rounded; glossy dark purplish-bronzy-fuscous: cilia concolorous. Hindwings and cilia dark fuscous; 2-4 remote, equidistant, 7 on lower surface with downwards-directed fringe of long hairs.

CAMEROONS, Lolodorf, February, March (*Conradt*); 2 ex. (Coll. Paravicini). The abnormal neuration of hindwings is found in two other species already admitted into the genus, the other characteristic features being present as usual; the curious fringe of hairs in hindwings is a quite unique structure.

*Macrobathra antimeloda*, n. sp.

♂. 15 mm. Head white. (Palpi missing). Thorax dark fuscous, tegulae mixed white. Abdomen ochreous-whitish. Forewings dark fuscous, bases of scales whitish, some scattered blackish scales; markings white; a narrow-triangular spot from costa before  $\frac{1}{4}$  reaching half across wing, an oblong spot on dorsum rather beyond this, a small spot on middle of costa, a moderately large hemispherical spot on dorsum slightly beyond this, a moderately large subtriangular spot on costa at  $\frac{4}{5}$ , and a small spot on termen beneath it: cilia whitish-grey, a few dark fuscous specks, and fuscous line before tips. Hindwings rather dark grey, cilia grey.

MADAGASCAR, Imerina (*Camboué*); 1 ex. (Coll. Paravicini). Probably allied to *distincta* Wals.

**Promalactis isothea, n. sp.**

$\delta \text{ } \varnothing$ . 11-13 mm. Head shining bronzy-grey. Palpi orange, terminal joint black. Antennæ black dotted white, apical fourth white except extreme tip, ciliations  $\delta$  nearly 2. Thorax shining dark bronzy-grey, tegulæ orange. Forewings orange; five bright iridescent-violet-metallic black-edged narrow fasciæ, first basal, edge oblique, second at  $\frac{1}{4}$ , dilated towards costa, third median, rather approximated and sometimes connected with second on dorsum, fourth interrupted, forming a dot on costa at  $\frac{2}{3}$  and a vertical streak from tornus, fifth terminal, extending round apex over apical fifth of costa, and joining fourth at tornus, between fourth and fifth a series of five or six fine black lines on veins: cilia dark grey, lighter towards base, on termen outer  $\frac{2}{5}$  pale yellowish. Hindwings dark fuscous; cilia rather dark grey.

MADRAS, Kodaikanal, Palnis, 7000 feet, August, "resting on leaves of *Poinsettia* on sunny afternoons" (Fletcher); 4 ex. At first sight extremely similar to *callimetalla* from the same region, but in that species the groundcolour of forewings is yellow with deep orange patches, the fourth fascia is connected with third by a median line and not interrupted, the black neural lines between fourth and fifth fasciæ are absent, and the terminal cilia are wholly orange.

**ALOMENARCHA, n. g.**

Head with appressed scales; ocelli posterior; tongue developed. Antennæ  $\frac{4}{5}$ , basal  $\frac{2}{3}$  rough-scaled dorsally,  $\delta$  more strongly and pectinated with long ciliated filaments, scape moderate, without pecten. Labial palpi long, recurved, second joint rather thickened with appressed scales, terminal joint as long as second, slender, acute. Maxillary palpi rudimentary. Posterior tibiæ rough-scaled above. Forewings 1 b furcate, 2 and 3 long-stalked from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings 1, trapezoidal-ovate, cilia  $\frac{1}{5}$ ; 2 from  $\frac{3}{5}$ , 3 and 4 connate or short-stalked, 5 straight, parallel, 6 and 7 slightly approximated towards base.

A development of *Dasysera*.

**Alomenarcha phalarista, n. sp.**

$\delta \text{ } \varnothing$ . 27 mm. Head, thorax dark fuscous, face and apical half of tegulæ ochreous-whitish. Palpi fuscous. Antennæ black, a white band towards apex. Abdomen orange-yellow, 4 apical segments blackish towards base. Forewings elongate, posteriorly dilated, costa anteriorly straight, somewhat bent at  $\frac{2}{3}$  and gently arched towards obtuse-pointed apex, termen obliquely rounded; blackish, irregularly sprinkled pale yellowish, beyond cell with fine irregular pale yellowish lines on veins; a longitudinal pale yellowish streak from base of costa to a spot in disc at  $\frac{1}{3}$ , and another subdorsal from base to middle or reaching a narrow irregular yellowish transverse fascia at  $\frac{2}{3}$  constricted in middle and not quite reaching margins; a yellowish marginal line round apical part of costa and termen: cilia dark fuscous. Hindwings

orange; ♂ basal third and broad terminal fascia occupying  $\frac{1}{3}$  at apex, ♀ base narrowly suffused blackish and moderately broad blackish terminal fascia, occupying nearly  $\frac{1}{4}$  at apex and narrowed to a point at tornus; cilia fuscous; a darker basal line.

SILHET, Chandkira, 1 ♂ (damaged) in my Coll.; SIKKIM, 1 ♀ (Coll. Paravicini).

### *Cyphothyris ophryodes* Meyr.

An example in fine condition from Hydrographer Mts., New Guinea (Coll. Paravicini) permits improvement of description and correction of inaccuracies; ♂ 14 mm., forewings white towards base, brown plical streak in this example absent, plical stigma little marked, preceded by a large tuft of scales, a rhomboidal fuscous blotch mixed blackish and whitish extending on costa from middle to  $\frac{3}{4}$  and reaching half across wing, followed by somewhat prominent costal scales and an angulated white line from costa to tornus, its angle projecting to touch termen in middle, apical area dark fuscous above this: hindwings 2 from well beyond middle of lower margin, curved, 3 and 4 connate, 5 parallel, 6 and 7 approximated towards base.

### *Allotalanta ockthotoma*, n. sp.

♂ ♀. 12 mm. Head whitish-ochreous. Palpi ochreous-whitish, terminal joint with dark fuscous supramedian ring. Thorax pale ochreous, shoulders with a dark fuscous mark. Forewings elongate, costa gently arched, apex pointed, termen faintly sinuate, oblique; pale ochreous-yellowish suffused white on margins of markings; an elongate blackish patch extending along costa from  $\frac{1}{3}$  to apex, cut by very oblique white strigæ from costa at middle and towards apex: a blackish patch occupying dorsal  $\frac{2}{3}$  from base to tornus, edge posteriorly irregular; one or two fine blackish dashes in disc toward apex: cilia ochreous-whitish, on costa blackish (imperfect). Hindwings grey, darker posteriorly, hyaline space large, elongate-oval; cilia light grey, at apex an ochreous-white suffused spot.

CAMEROONS, Lolodorf and Johann-Albrechts Höhe Station (Conradt); 2 ex. (Coll. Paravicini).

### *Allotalanta tephroclystis*, n. sp.

♂. 15 mm. Head light brown, face whitish-ochreous. Palpi whitish, second joint with fuscous lateral streak on basal half, terminal joint with fine basal and median fuscous rings. Thorax brownish, anterior margin suffused dark purplish-fuscous. Forewings elongate, costa slightly arched, more strongly towards pointed apex, termen slightly sinuate, oblique; light pinkish-brown; an irregular undefined streak of dark grey suffusion along costa from  $\frac{1}{5}$  to apex, interrupted about  $\frac{4}{5}$ ; discal stigmata grey, surrounded by pale rings of groundcolour, adjoining this grey suffusion; some grey suffusion about upper part of termen: cilia

whitish-ochreous, with grey subbasal and subapical lines, becoming darker and confluent on costa. Hindwings grey; cilia light grey, a darker basal shade.

CAMEROONS, Johann-Albrechts Höhe Station (*Conradt*); 1 ex. (Coll. Paravicini).

**Piloprepes phthoromima, n. sp.**

♂. 18 mm. Head white, crown centrally suffused dark grey, face mixed grey and black. Palpi white, second joint suffused grey on basal  $\frac{2}{3}$  and with a blackish subapical ring; terminal joint mixed dark grey. Antennal ciliations fasciculated (4). Thorax white mixed grey and dark brownish. Forewings suboblong, costa anteriorly moderately arched, somewhat sinuate beyond middle, edge broadly prominent with rough projecting scales before this, apex obtuse, termen obliquely rounded; white, irregularly mixed or sprinkled grey; a very undefined basal patch of grey and dark grey suffusion, including an ochreous-brown suffused spot; an oblique suffused grey streak in disc before middle, and one from posterior edge of costal scale-prominence: cilia long, white, basal half suffusedly mixed grey. Hindwings white; two or three short streaks of grey suffusion between veins towards apex; cilia grey, round apex and upper part of termen whitish on outer  $\frac{2}{3}$ .

BRITISH NEW GUINEA, Owgarra (*Meek*); 1 ex. (Coll. Paravicini). A curious form.

**PYROPHRACTIS, n. g.**

Head with appressed scales, sidetufts loosely raised; ocelli posterior; tongue developed. Antennæ  $\frac{3}{4}$ , ♀ shortly ciliated (♂ probably evenly ciliated), scape elongate, with developed pecten. Labial palpi very long, recurved, second joint with appressed scales, terminal joint as long as second, somewhat roughened with scales anteriorly, acute. Maxillary palpi very short, filiform, appressed to tongue. Posterior tibiæ with rough scales above. Forewings 1 b furcate, 2 from angle, 7 and 8 stalked, 7 to apex, 11 from middle. Hindwings 1, trapezoidal-ovate, cilia  $\frac{2}{5}$ ; 3 and 4 connate, 5 rather curved and approximated to 4 at base, 6 and 7 parallel.

Allied to *Machimia*; general characters not very pronounced, but will not fall within any other genus.

**Pyrophractis isotherma, n. sp.**

♀. 20 mm. Head, thorax red mixed yellow. Palpi crimson irrorated dark fuscous, extreme tip of second joint, and basal, median, and apical rings of terminal pale yellow. Abdomen whitish-ochreous. Forewings moderate, rather dilated posteriorly, costa slightly arched, apex obtuse, termen rounded, little oblique; red, veins irregularly lined light ochreous-yellow; three slightly curved oblique dark purple-fuscous lines, first from base of costa to near base of dorsum, somewhat toothed, second from costa at  $\frac{1}{3}$  to middle of dorsum, third from costa at  $\frac{3}{5}$  to tornus, a branch

running from upper part of this to costa before apex: cilia light purplish-grey mixed dark grey, outer half suffused ochreous-whitish on costa and termen (leaving an apical bar), tornal area wholly greyish. Hindwings whitish-rose-pink, basal hairs ochreous-whitish; cilia ochreous-whitish.

BRITISH NEW GUINEA, Aroa R. (*Meek*); 1 ex. (Coll. Paravicini).

### EPIMORYCTIS, n. g.

Head loosely scaled, sidetufts roughly spreading; ocelli posterior; tongue developed. Antennæ  $\frac{3}{4}$ , ♂ strongly ciliated, scape elongate, stout, without pecten. Labial palpi very long, recurved, second joint thickened with dense scales, somewhat rough beneath, terminal joint as long as second, moderate, acute. Maxillary palpi very short, loosely scaled, appressed. Posterior tibiæ with appressed scales, rough-haired beneath on apical half. Forewings 1b furcate, 2 from towards angle, 3-5 approximated, 7 absent, 8 and 9 stalked, 11 from middle. Hindwings 1, elongate-ovate, cilia  $\frac{2}{5}$ ; 2 from  $\frac{3}{4}$ , 3 and 4 connate, 5 slightly curved, 6 and 7 slightly approximated.

A development of *Machimia*.

#### *Epimoryctis percnorma*, n. sp.

♂. 21 mm. Head whitish-ochreous partially mixed fuscous. Palpi ochreous-whitish, second joint with dark fuscous lateral streak. Antennal ciliations  $2\frac{1}{2}$ . Thorax pale ochreous (discoloured). Forewings rather elongate, slightly dilated, costa gently arched, apex obtuse, termen rounded, somewhat oblique; pale ochreous; costal edge whitish-ochreous, beneath this some brownish suffusion and dark fuscous irroration throughout, continued along termen to tornus, a similar spot on fold near base; stigmata blackish, plical obliquely beyond first discal, a fine incomplete line of blackish scales between discal; an angulated transverse band of brownish suffusion and dark fuscous sprinkling beyond cell, obsolete towards dorsum and interrupted above middle; a series of small blackish spots near and parallel to posterior third of costa and termen; marginal suffusion round apical part of costa and termen tending to be cut by pale lines on veins: cilia whitish-ochreous, an interrupted dark fuscous subbasal line. Hindwings light grey; cilia grey-whitish, round apex a grey antemedian line.

BRAZIL, Serra de Communaty, Pernambuco; 1 ex. (Coll. Paravicini).

#### *Machimia elaphodes*, n. sp.

♂. 23 mm. Head, thorax pale brownish-ochreous. Palpi whitish, second joint dark fuscous towards base. Antennal ciliations 3. Forewings suboblong, slightly narrowed anteriorly, costa gently arched, apex obtuse, termen slightly rounded, little oblique; light brownish-ochreous, a few minute fuscous specks; extreme costal edge blackish near base; plical and second discal stigmata very small, dark grey, first discal forming a conspicuous small roundish blackish-grey spot; a series of small grey dots from beneath costa at  $\frac{2}{3}$  very obliquely outwards, right-angled opposite apex and then

rather excurved to dorsum at  $\frac{2}{3}$ : cilia pale brownish-ochreous. Hindwings and cilia ochreous-grey-whitish.

BOLIVIA, Cochabamba (*Germain*); 1 ex. (Coll. Paravicini). Perhaps nearest *pagana*.

### *Syscalma prymnæa*, n. sp.

Of two other examples seen, one from the Hydrographer Mts. (certainly only a variety) has the forewings clear orange except a small basal patch and moderately broad terminal fascia attenuated to tornus.

### *Barantola panarista* Turn.

An example from Biagi, Mambare R., New Guinea (Coll. Paravicini), differs from the description of the unique Queensland type as follows: ♂ 16 mm. (Turner's 25 mm. for ♀ is perhaps a misprint for 15), crimson tinge absent except in apical edge of forewings, costal streak absent on basal fifth, hindwings without grey apical suffusion; these differences may be individual, sexual, or geographical, but not specific.

### *Barantola pulcherrima* Walk.

A ♂ from Biagi, Mambare R., New Guinea, has the collar white edged red, the antemedian fascia of forewings represented only by a triangular white red-edged projection from costal streak, a black basal line in cilia round costa, apex, and upper part of termen. The species seems to be variable, no two of the few examples I have seen being alike, and this form is probably not specifically distinct.

### *Scorpiopsis exanthistis*, n. sp.

♂. 14 mm. Head, thorax pale yellowish, obscurely lined dull crimson suffusion. Palpi deep ferruginous, tip white. Forewings somewhat elongate, posteriorly dilated, costa gently arched, apex rounded-obtuse, termen nearly straight, slightly oblique; 3 and 4 out of 2; pale yellow, reticulated throughout orange-red; a fuscous costal streak from base to middle, costal edge of this blackish with a white mark before middle; a spot of fuscous suffusion on end of cell, whence a vague streak of suffusion runs to a fuscous mark on costa at  $\frac{3}{4}$ ; a fuscous terminal streak including some yellowish dots: cilia yellow, some slight reddish bars near base. Hindwings and cilia pale yellowish.

BRITISH NEW GUINEA, Hydrographer Mts., 2500 feet (*Eichhorn*); 1 ex. (Coll. Paravicini). The other three species of the genus have veins 2-4 of forewings approximated only, but a new genus is unnecessary here, the structure and habit being otherwise similar, and my published key to the genera needing no modification.

### *Scorpiopsis diplaneta*, n. sp.

♂. 18-19 mm. Head crimson somewhat mixed pale yellow. Palpi crimson, anterior edge of second joint fuscous or white,

a subapical ring of second joint, and basal and apical bands of terminal joint pale yellow, thickened scales of terminal joint partly infuscated. Thorax crimson, dorsally somewhat infuscated, four anterior pale yellow spots and three posterior dots more or less developed. Forewings moderately broad, costa rather strongly arched, apex rounded, termen rounded, little oblique; 7 to costa; crimson, variably more or less suffused fuscous except on margins of markings, with numerous small pale yellow spots arranged in series; extreme costal edge white except towards extremities; a round snow-white spot in disc at  $\frac{2}{3}$ , and smaller one obliquely beyond and above it: cilia light yellow, base suffused crimson, towards tornus suffused light grey. Hindwings ochreous-whitish, faintly or sometimes strongly suffused rose-pink; apex sometimes narrowly grey; cilia ochreous-whitish.

BRITISH NEW GUINEA, Biagi, Mambare R., 5000 feet, and Kumusi R. (Meek); DUTCH NEW GUINEA, Oetakwa R., Snow Mts.; 3 ex. (Coll. Paravicini).

#### *Scorpiopsis rhodoglaaca*, n. sp.

♂. 23 mm. Head crimson, several small whitish spots and marks. (Palpi broken.) Thorax crimson, four small anterior pale yellow spots, and four grey spots behind these. Forewings rather broad, costa strongly arched, apex rounded, termen rounded, nearly vertical; crimson, with numerous small pale yellow spots arranged in series; costal edge crimson, beneath this a narrow dark grey streak throughout continued round termen to tornus; a small dark grey spot on dorsum at  $\frac{2}{5}$ , from which some slight grey suffusion runs to dark grey second discal stigma; a curved fascia of grey suffusion tending to be broken into streaks from  $\frac{2}{3}$  of costa to dorsum before tornus; cilia pale yellow, a crimson basal line, tornal area light grey. Hindwings whitish, suffused light rosy-pink except towards base, basal hairs tinged ochreous; cilia yellow-whitish.

QUEENSLAND, Kuranda (Dodd); 1 ex. (Coll. Paravicini).

#### *Octasphales niphadosticha*, n. sp.

♀. 17 mm. Head grey, face white. Palpi second joint grey, apex white, terminal joint white with two dark grey bands. Thorax grey, shoulders white. Forewings oblong, costa anteriorly moderately arched, with antemedian projection of rough scales, appearing faintly sinuate between this and abrupt origin of cilia, apex rounded, termen rounded, vertical; grey, darker on veins, strewn throughout with numerous white dots forming longitudinal series, posteriorly on veins; costal edge white between antemedian scale-projection and cilia: cilia white, basal half barred grey. Hindwings and cilia grey.

BRITISH NEW GUINEA, Kumusi R. (Meek); 1 ex. (Coll. Paravicini). The costal scale-projection suggests resemblance with *Acria* and *Piloprepes*, perhaps mimetic.

*Depressaria nyctalopis*, n. sp.

♀. 23 mm. Head ochreous-whitish sprinkled light grey, side-tufts mixed dark fuscous. Palpi ochreous-whitish, second joint irrorated dark fuscous on basal half, tip rosy-tinged above a dark fuscous subapical ring, terminal joint with subbasal and supramedian dark fuscous bands. Thorax ochreous-whitish irrorated brownish and dark fuscous. Forewings elongate, costa gently arched, apex rounded, termen rather obliquely rounded; 2 and 3 stalked; whitish-ochreous irregularly strigulated brownish, a few dark fuscous scales; costal edge tinged rosy from base to beyond middle, with small black spots or dots throughout; minute black dots near base above middle and above dorsum; a blotch of grey suffusion and irroration on dorsum at  $\frac{1}{3}$ ; a lunulate black mark in disc at  $\frac{1}{3}$ , preceded by some brown suffusion and black irroration, and followed first by some whitish suffusion, then by a small blackish spot, close beyond which is a larger trapezoidal black spot; a fine black dash towards costa at  $\frac{2}{3}$ ; some slight indistinct dark fuscous terminal marks: cilia whitish-ochreous, a brownish subbasal shade. Hindwings light greyish; cilia whitish, a light grey subbasal shade.

COMORO Is., Grand Comoro; 1 ex. (Coll. Paravicini). Perhaps nearest *dryocrates*.

*Hypercallia deflorescens*, n. sp.

♂ ♀. 18-21 mm. Head whitish-ochreous, sides of crown brownish or fuscous, face ♂ fuscous. Palpi whitish, second joint fuscous except apex, scales roughly expanded towards apex above, terminal joint with fuscous median band, hardly perceptibly thickened. Antennal ciliations ♂ long on basal half. Thorax light brownish-ochreous or greyish-ochreous, ♂ infuscated. Forewings suboblong, costa gently arched, apex obtuse, termen slightly sinuate, rather oblique; 9 separate; pale greyish-ochreous irregularly sprinkled fuscous and dark fuscous, costal half suffused ochreous-fuscous, veins sometimes marked brown streaks; discal stigmata dark fuscous, first followed by a white dot (sometimes obsolete) and then by a short reddish dash; a blackish-grey elongate mark on dorsum towards tornus; a terminal series of small blackish dots: cilia light brownish. Hindwings ochreous-grey-whitish or whitish-grey; cilia whitish or ochreous-whitish, sometimes a light greyish-ochreous subbasal shade.

SIKKIM, Gangtok, 6000 feet, April, May (Colonel F. M. Bailey): 7 ex. A variable and obscure form, but interesting geographically.

*Diocosma rhodopola*, n. sp.

♂. 15 mm. Head white. Palpi white, second joint grey externally on basal half, extreme base of terminal joint dark grey. Thorax fuscous, slightly purple-tinged anteriorly. Forewings rather elongate, costa gently arched, apex rounded-obtuse, termen obliquely rounded; light violet-grey, dorsal half suffused darker grey; a suffused rose-pink costal streak from base to apex: cilia

rosy-whitish, on costa rosy, on tornus grey. Hindwings light grey; cilia grey-whitish.

KENIA, Kibwezi-Uganda; 1 ex. (Coll. Paravicini).

### *Calliphractis phyllograpta* Meyr.

Having now seen a ♂ (from Cameroons, in Coll. Paravicini) I find that vein 11 of forewings is absent in that sex, just as in *Orygocera*, of which genus it must be considered a development; terminal joint of palpi with basal and median blackish rings: thorax with green posterior spot; some slight variations of detail in markings of forewings, otherwise essentially as in ♀.

### *Orygocera liliacea*, n. sp.

♂. 14 mm. Head, palpi, thorax white sprinkled dark grey. Forewings rather elongate, somewhat dilated, costa gently arched, apex obtuse-pointed, termen nearly straight, somewhat oblique; pale whity-greenish; costal edge dark fuscous on basal third; three shining white blotches edged dark fuscous, first hemispherical, on dorsum about  $\frac{1}{3}$ , second fasciate, inwards-oblique, from dorsum about  $\frac{3}{4}$  reaching half across wing, third roundish, on costa beyond middle, unmargined posteriorly, touching second; wing beyond these wholly suffused white except towards tornus, slightly sprinkled fuscous; a terminal series of fuscous specks: cilia white, outer half grey speckled darker. Hindwings and cilia light grey.

CAMEROONS, Lolodorf, February (Conradt); 1 ex. (Coll. Paravicini).

### *Orygocera amphichelota*, n. sp.

♂. 16 mm. Head whitish, crown suffusedly irrorated grey. Palpi white, irrorated grey except towards base. Thorax white irrorated dark grey. Forewings rather elongate, costa gently arched, apex obtuse-pointed, termen hardly sinuate, rather oblique; white, rather thinly sprinkled brownish or fuscous, more densely towards dorsum; costa towards base suffused fuscous irroration; a dark fuscous streak along dorsum from middle to  $\frac{4}{5}$ , with a rather inwards-oblique slightly curved projection from anterior end reaching fold, and a shorter outwards-oblique projection from posterior end; cloudy rather dark fuscous dots on both angles of cell; a marginal series of small dark fuscous dots round apex and termen; cilia white, outer half grey-whitish, grey median and subapical and dark fuscous apical lines. Hindwings pale grey; cilia grey-whitish.

CAMEROONS, Johann-Albrechts Höhe Station (Conradt); 1 ex. (Coll. Paravicini).

### *Orygocera (?) magdalena*, n. sp.

♀. 23 mm. Head white. Palpi light pinkish sprinkled grey. Thorax whitish, tegulae tinged pink and sprinkled grey. Forewings rather elongate, costa gently arched, apex obtuse, termen nearly straight, little oblique; pale rose-pink, bases of scales pearly whitish, some scattered dark fuscous scales on posterior half; costal

edge white from before middle to apex ; a triangular rosy-grey blotch on dorsum beyond middle, partially suffused dark red-brown on margins and edged whitish ; a slender inwards-oblique reddish-brown streak from dorsum before tornus, edged white anteriorly, discal area tinged grey above this : cilia ochreous-whitish with two faint rosy lines. Hindwings pale grey with faint pinkish tinge, becoming grey-whitish anteriorly ; cilia ochreous-whitish with faint rosy tinge.

MADAGASCAR, Forest of Antsianaka (*Humblot*) ; 1 ex. (Coll. Paravicini). In absence of ♂ this might possibly be an *EpiphRACTIS*.

#### *Meleonoma nephospora*, n. sp.

♂. 14 mm. Head, thorax pale greyish-ochreous somewhat sprinkled fuscous, tegulae suffused fuscous. Palpi whitish, second joint irregularly speckled blackish, with blackish apical ring, terminal joint with slender blackish median ring. Antennæ ochreous-whitish ringed dark fuscous. Forewings elongate, costa gently arched, apex pointed, termen very obliquely rounded ; pale ochreous, thinly speckled fuscous ; an undefined patch of dark grey suffusion occupying costal half from base to  $\frac{2}{5}$  ; a semi-oval dark grey blotch on middle of costa ; stigmata blackish, first discal and plical transversely placed on edge of basal patch, second discal transverse ; an irregular terminal fascia of dark grey suffusion : cilia light grey speckled ochreous-whitish. Hindwings rather dark grey ; cilia grey.

SIKKIM, Gangtok, 6000 feet, May (Colonel F. M. Bailey) ; 1 ex. A second example, probably conspecific, has the groundcolour of forewings wholly suffused light grey and irrorated dark grey.

#### *Periacma delophanta*, n. sp.

♀. 17 mm. Head yellowish-white. Palpi whitish, second joint with dark fuscous apical band. Thorax dark fuscous, posterior extremity white. Forewings moderate, slightly dilated, costa gently arched, apex pointed, termen slightly sinuate, oblique ; dark fuscous ; stigmata obscure, cloudy, blackish, first discal edged white posteriorly, plical rather obliquely beyond first discal, indenting edge of a hemispherical white blotch on dorsum beyond middle ; a triangular white blotch on costa about  $\frac{3}{4}$  ; small whitish spots on termen beneath apex and at tornus : cilia grey, base whitish (imperfect). Hindwings grey ; cilia light grey.

MADRAS, Kodaikanal, Palnis, 7000 feet, August (Fletcher) ; 1 ex. Allied to *conophanta*.

#### *CRITOXENA*, n. g.

Head with appressed scales, sidetufts loosely spreading ; ocelli posterior ; tongue developed. Antennæ 1, scape moderate, with pecten. Labial palpi rather long, recurved, second joint with appressed scales, terminal joint rather shorter than second, slender, acute. Maxillary palpi rudimentary. Posterior tibiae clothed with long rough hairs above. Forewings 1 & 2 furcate, 3 very short,

stalked from angle, 7 and 8 stalked, 7 to costa, 11 from middle. Hindwings under 1, ovate-lanceolate, cilia 1; 2 from  $\frac{3}{4}$ , 3 and 4 connate, 5 and 6 stalked, 7 parallel.

Allied to *Scalideutis*.

### *Critoxena agraphopis*, n. sp.

♀. 17 mm. Head, palpi, thorax light greyish-ochreous. Forewings elongate, costa gently arched, apex pointed, termen extremely obliquely rounded; light greyish-ochreous; plical stigma indicated by one or two slightly raised fuscous scales, second discal hardly darker than groundcolour: cilia pale greyish-ochreous. Hindwings pale grey; cilia pale greyish-ochreous.

INDIA CENTRAL PROVINCES, Dehra Dun, 2000 feet, July (*Fletcher*); 1 ex.

## LAMPRONIADÆ.

### *CATOLBISTIS*, n. g.

Head rough-haired; ocelli posterior; tongue absent. Antennæ  $\frac{1}{2}$ , slender, joints elongate, ♂ ciliated, stalk rough-scaled above at base, scape moderate, without pecten. Labial palpi moderately long, subascending, clothed with loose long hairscales, projecting beneath on second joint and above on terminal, terminal joint as long as second, obtuse. Maxillary palpi rudimentary. Posterior tibiæ with appressed scales. Forewings 2 from towards angle, 7 and 8 coincident, to apex, 9 and 10 from towards angle, 11 from before middle. Hindwings 1, subovate, cilia  $\frac{3}{5}$ ; 2 from  $\frac{2}{3}$ , 3-7 nearly equidistant.

### *Catolbistis thrasymedes*, n. sp.

♂. 14 mm. Head, palpi ochreous-yellowish. Antennæ yellow-whitish ringed grey, ciliations  $1\frac{1}{2}$ . Thorax dark purple-grey. Forewings rather elongate, posteriorly considerably dilated, costa moderately arched, apex obtuse, termen obliquely rounded; dull purple suffusedly irrorated grey; a rather broad slightly curved brassy-yellowish fascia before middle: cilia grey. Hindwings and cilia grey.

UGANDA, Elgor, 11000 feet, August (*G. L. R. Hancock*); 1 ex. (Brit. Mus.).

## ADELIDÆ.

### *Nemotois xanthophracta*, n. sp.

♂. 15 mm. Head yellow-ochreous, face brassy-metallic, eyes large, not nearly approximated above. Palpi short, second joint grey with short projecting pale ochreous hairs, terminal joint pale ochreous. Antennæ slender, whitish, basal third grey. Thorax purplish-bronzy-metallic. Forewings rather short, moderately

dilated, costa bent about  $\frac{2}{3}$ , apex obtuse-pointed, termen obliquely rounded; bright deep coppery-purple; an orange-yellow blackish-edged basal patch occupying  $\frac{1}{4}$  of wing, edge angulated and interrupted in middle by a lobe of groundcolour extending inwards to near base of costa; two rather narrow orange-yellow blackish-edged fasciæ, first slightly excurved, midway between basal patch and second, second straight, median; a broad fascia of black irroration from  $\frac{2}{3}$  of costa to tornus: cilia dark purple-grey. Hindwings dark grey; cilia grey. Posterior tibiæ with fringe of very long fine hairs above.

BOMBAY, Mahabaleshwar, October (*Maxwell*); 1 ex. Nearest *athlophora*.

#### *Nemotois ischnodesma* Meyr.

1 ♂, Pidaung, Burma, April. Antennæ whitish ringed dark fuscous, towards base dark fuscous; posterior tibiæ with fringe of long whitish hairs.

#### *Nemotois porphyraspis*, n. sp.

♀. 16 mm. Head grey, face bright brassy-metallic. Palpi short, whitish. Antennæ (broken) white, basal half (?) thickened with dark purple-bronzy scales. Thorax shining brassy-bronze. Forewings subtriangular, costa strongly arched posteriorly, termen oblique; bright coppery-golden, becoming bright brassy-bronze towards base; a broad deep purple postmedian fascia, strewn light ochreous scales, only moderately broad in middle but strongly dilated towards both extremities, and a similar round apical blotch: cilia purplish, suffused coppery-golden towards base. Hindwings and cilia dark fuscous.

KENYA, Nairobi, Oct. (*Dr. G. A. K. Marshall*); 1 ex. (Brit. Mus.). Belongs to group of *fasciella*.

### COLEOPHORIDÆ.

#### *Coleophora summivola*, n. sp.

♂. 17 mm. Head grey-whitish. Palpi light grey, base white, second joint with short apical scale-projection, terminal joint little over half second. Antennæ grey-whitish. Forewings whitish-ochreous, costa slenderly white from base to  $\frac{2}{5}$ , some scattered grey scales along veins: cilia ochreous-whitish, tinged grey on basal half. Hindwings and cilia pale grey.

TIBET, Gyantse, 14500 feet, July (*Colonel F. M. Bailey*); 1 ex. Perhaps allied to *chordoscelis*.

#### *Coleophora nivifera*, n. sp.

♂. 13 mm. Head, thorax yellow-whitish. Palpi white, second joint tinged grey-yellowish externally, without projection, terminal joint  $\frac{2}{3}$ . Antennæ rather stout towards base, white. Forewings

white, tinged grey-yellowish suffusion between costal veins; a short fine grey line on costal edge towards  $\frac{3}{4}$ : cilia white. (Hindwings missing.)

TIBET, Yatung, 10000 feet (*Colonel F. M. Bailey*); 1 ex.

### EPERMENIADÆ.

#### *Epermenia sublimicola*, n. sp.

♂ ♀. 17-19 mm. Head, palpi, thorax whitish speckled grey. Forewings elongate-lanceolate; white, speckled grey and dark grey; discal stigmata small, black, widely remote, second sublinear, a faint streak of pale brownish suffusion between and beyond these, and another interrupted along fold; a small black scaletooth on dorsum beneath first discal, some slight blackish irroration indicating a short oblique streak from this; a second small dorsal scaletooth with three or four black scales midway between stigmata; a third somewhat before second discal, and a minute one beyond it; a minute black dot at apex, and one or two on costa before it: cilia light grey, at and beneath apex some darker lines and dark grey basal speckling. Hindwings and cilia grey.

TIBET, Gyantse, 13000 feet (*Colonel F. M. Bailey*); 2 ex.  
Allied to *macescens*.

#### *Epermenia parasitica*, n. sp.

♂. 8-9 mm. Head, thorax whitish closely irrorated blackish-grey. Palpi dark grey, apex of joints suffused whitish. Forewings rather narrow, parallel-sided, moderately pointed, dorsum with rough scales forming three or four undefined projections; grey-whitish, apex of scales dark grey, forming a close rather irregular striolation, more or less largely suffused ochreous-brown except anteriorly; three small black dots in a longitudinal row in disc from  $\frac{1}{4}$  to  $\frac{3}{4}$ ; several dark grey transverse spots from costa; a black apical dot: cilia light grey, with rows of blackish points. Hindwings grey; cilia light grey.

JAVA, Buitenzorg, bred from *Loranthus* (*Dr. S. Leefmans*); 2 ex. (type Brit. Mus.). Allied to *dracontias*.

## INDEX.

---

- absolutrix, 547.  
abstrusa, 427.  
accincta, 5.  
acharnaea, 348.  
acherusia, 29.  
achyritis, 257.  
aciculata, 468.  
aemophanes, 409.  
acontarcha, 295.  
acratopa, 288.  
Acrocercops, 85, 292,  
  407, 580.  
Acroclita, 140, 333,  
  430, 598.  
acroglypta, 505.  
Acrolepia, 359.  
Acrolophus, 76, 329,  
  552.  
Acroplectis, 370.  
actinopa, 481.  
acuta, 354.  
Adaina, 568.  
Adela, 572.  
adoratrix, 200.  
Adoxophyes, 453.  
adulans, 220.  
adytodes, 196.  
aedificata, 507.  
Aegeria, 266, 371, 465,  
  584.  
aenictopa, 483.  
Aeolanthes, 146.  
aeolodoxa, 423.  
aeolornis, 605.  
aeolozona, 252.  
Aeonoxena, 431.  
aericincta, 466.  
aerinotata Butl., 192.  
aethalopa, 102.  
aetherias, 361.  
affusa, 40.  
agalmatopa, 305.  
Agathactis, 501.  
  
agatharcha, 253.  
Agdistis, 95.  
aggesta, 147.  
aglaocarpa, 113.  
aglaopa, 389.  
aglaospila, 463.  
aglaoxantha, 116.  
aglypta, 174.  
Agonoxena, 84.  
agraphopis, 624.  
agraria, 210.  
Agrioceros, 417.  
agriodes, 350.  
Agriophanes, 600.  
Agriophara, 560.  
agrioschista, 365.  
agrippina, 594.  
agyrtodes, 344.  
albifrons Z., 557.  
albiramis, 51.  
albitibiana Snell., 447.  
alcea, 72.  
alcestis, 40.  
alcmaeonis, 455.  
alcyonopa, 301.  
aleatoria, 461.  
alicia, 461.  
alliciens, 305.  
alligans Butl., 192.  
Allotalanta, 310, 381,  
  616.  
allotriopa, 58.  
allutella Reb., 552.  
alluvialis, 204.  
Alomenarcha, 615.  
alopecopa, 518.  
altivaga, 137.  
altivola, 486.  
Alucita, 299, 567.  
amauropis, 1.  
amblycryptis, 516.  
amnesta, 36.  
Amorbia, 260.  
  
ampelurga, 146.  
ampherista, 163.  
amphichelota, 622.  
amphichlora, 4.  
amphicroca, 151.  
amphignosta, 244.  
amphilecta, 145.  
amphilocha, 128.  
amphimnesta, 436.  
Amphithales, 268.  
Ampithera, 587.  
amphitrocta, 322.  
amphixysta, 500.  
Amphixystis, 399.  
amplificata, 153.  
Anacampsis, 281, 507.  
Anacathartis, 383.  
anachasta, 467.  
anamnestis, 450.  
anaphanta, 180.  
Anaphantis, 593.  
Anarsia, 281, 353, 512.  
Anastathma, 266.  
Anathamna, 441.  
ancalodes, 395.  
anceps Butl., 192.  
Anchicremna, 246.  
Anchinia, 100.  
ancylacma, 205.  
Ancylis, 438.  
Ancylometis, 51.  
anelicta, 312.  
anemolia, 533.  
anisectis, 19.  
anisochrysa, 415.  
anisodora, 98.  
anisogramma, 353.  
anisota, 374.  
annuligera, 397.  
anomogramma, 549.  
Anomologa, 308, 533.  
anomorcta, 277.  
Anomoxena, 5.

*Antaeotricha*, 162.  
*Anthistarcha*, 484.  
*anthistis*, 508.  
*anthophaga*, 392.  
*anthophanes*, 438.  
*anthostoma*, 441.  
*anthracaula*, 512.  
*anthracuris*, 295.  
*antiastis*, 514.  
*anticeros*, 422.  
*anticipans*, 342.  
*anticoma*, 536.  
*Anticrates*, 121, 361,  
 594.  
*Antictenista*, 337.  
*antigama*, 270.  
*Antigambra*, 321.  
*Antigraptis*, 613.  
*antilyra*, 215.  
*antimeloda*, 614.  
*antiphanova*, 372.  
*Antispastis*, 307.  
*Antispila*, 261, 541.  
*antisticha*, 285.  
*antisticta*, 511.  
*antitacta*, 219.  
*antitona*, 364.  
*Apatetris*, 51, 480.  
*Apethistis*, 291.  
*aphanozona*, 290, 356.  
*aphromorpha*, 7.  
*aphroneura*, 572.  
*apochranta*, 609.  
*apoplectica*, 594.  
*aporopis*, 477.  
*arabarcha*, 572.  
*arachnogramma*, 252.  
*archiphenax*, 404.  
*Ardeutica*, 260.  
*argicerauna*, 221.  
*argo*, 540.  
*argocymba*, 232.  
*argonais*, 224.  
*argutus*, 299.  
*argyRACTIS*, 8.  
*Argyroploce*, 68, 143,  
 339, 444, 601.  
*argyrospila*, 52.  
*Argyrotoxa*, 461, 612.  
*aristarcha*, 262.  
*aristopa*, 134.  
*Aristotelia*, 6, 272, 343,  
 481.  
*armigera*, 60.  
*Arctromima*, 532.  
*arquata*, 307.  
*Articolla*, 601.  
*articulata*, 274.  
*arvicola*, 70.

*Asapharcha*, 555.  
*asbolodes*, 349.  
*asbolopa*, 51.  
*asbolopis*, 403.  
*Ascalenia*, 60.  
*ascetica*, 311.  
*Aschistophleps*, 585.  
*asmenopa*, 133.  
*asphalopis*, 188.  
*associata*, 214.  
*astaurota*, 580.  
*asterodoxa*, 462.  
*asthenodes*, 6.  
*astigmatica*, 467.  
*astiopa*, 581.  
*Astoxena*, 593.  
*astragalopa*, 327.  
*astrocosma*, 136.  
*astyarcha*, 584.  
*atacta*, 343.  
*Atelosticha*, 472.  
*Athlostola*, 97.  
*Athrinacia*, 518.  
*athyris*, 415.  
*atlantis*, 85.  
*atmastra*, 506.  
*atmochlora*, 603.  
*atmospora*, 209.  
*atomophora*, 429.  
*Atoponeura*, 270.  
*atricapsis*, 602.  
*atriflora*, 609.  
*atrinotata*, 542.  
*Atteva*, 415, 419.  
*Atychia*, 302.  
*aucupatrix*, 499.  
*augescens*, 206.  
*aulacopis*, 6.  
*aulaeodes*, 254.  
*aularia*, 66.  
*aulodoma*, 77.  
*autampyx*, 161.  
*autocosma*, 124.  
*autocrena*, 575.  
*autodyas*, 525.  
*Autosticha*, 50, 532.  
*auxoleuca*, 223.  
*auxoplaca*, 255.  
*auxoptila*, 276.  
*Axyrostola*, 29.  
*axysta*, 363.  
  
*baccheutis*, 109.  
*Bactra*, 442.  
*bactropa*, 336.  
*Baeonoma*, 160.  
*baeopis*, 523.  
*Balanoptica*, 592.  
*balanospila*, 555.  
  
*baliochlora*, 535.  
*baliomicta*, 424.  
*balsamopa*, 16.  
*banausopa*, 526.  
*Barantola*, 619.  
*barbara*, 92.  
*barydelta*, 15.  
*barysphena*, 34.  
*barysticta*, 322.  
*barythyma*, 530.  
*batanella Busck*, 494.  
*bathroptila*, 517.  
*bathrotoma*, 189.  
*Batrachedra*, 92, 394,  
 547.  
*Battaristis*, 505.  
*Bedellia*, 397.  
*Beltheca*, 270.  
*beryllina*, 335.  
*Besciva*, 270.  
*biannulella Chamb.*,  
 352.  
*biarcuata*, 238.  
*bidens*, 564.  
*bidentifera*, 458.  
*binigrata*, 456.  
*binocularis*, 484.  
*binummulata*, 513.  
*biremis*, 256.  
*bispiculata*, 23.  
*Blastobasis*, 145.  
*bolistis*, 203.  
*Bondia*, 589.  
*Borkhausenia*, 381,  
 467, 574.  
*bosquella Chamb.*, 462.  
*bovina*, 428.  
*Brachmia*, 46, 290,  
 356, 527.  
*Brachyacma*, 510.  
*brachyclista*, 434.  
*brachyplaca*, 228.  
*brachyplecta*, 412.  
*brackenridgella Busck*,  
 493.  
*brevisecta*, 601.  
*britomartis*, 126.  
*bryianthes*, 245.  
*Bryonympha*, 560.  
*bryoscopa*, 445.  
*Bucculatrix*, 265, 398.  
*Bucolarcha*, 515.  
*bulligera*, 459.  
*byblinopa*, 154.  
*byssina*, Zell., 192.  
*byssophanes*, 230.  
  
*Cacoecia*, 53, 106, 455,  
 614.

- caecivaga, 480.  
 caenochytis, 187.  
 caerula, 323.  
 calais, 570.  
 calamitosa, 563.  
 Calamotypa, 272.  
 calcarata, 117.  
 calculatrix, 7.  
 calens, 9.  
 callichrysa, 266.  
 callicirrha, 86.  
*Calliphractis*, 476, 622.  
*Calliprora*, 281, 502.  
 callipyra, 416.  
 callirrhoda, 9.  
 callisarca, 105.  
*Callithrinca*, 361.  
 columnians, 284.  
 calycophthalma, 293.  
*Camineutis*, 544.  
*camptospila*, 202.  
*Campstrodoxa*, 144.  
 cantatrix, 221.  
 cantharites, 464.  
 canthonias, 438.  
 capnobathra, 605.  
 capnocrossa, 197.  
 capnographa, 402.  
*Capnolocha*, 152.  
 captatrix, 126.  
*Capua*, 106, 247, 369,  
     452.  
 carabopa, 129.  
 carcerata, 339.  
 carcinopis, 522.  
*Carmentina*, 597.  
*Carposina*, 137, 402,  
     540.  
 carycodes, 256.  
 caryodesma, 184.  
*Cassiopeia*, 362.  
 cassiteris, 585.  
 cassiterota, 40.  
 castalia, 561.  
 castigata, 318.  
 cataclina, 44.  
 catacrossa, 347.  
 catagrapta, 332.  
*Catalectis*, 64, 82.  
 catalina, 267.  
 catapeltica, 119.  
 catectis, 49.  
 catericta, 354.  
 cathagnista, 165.  
 catharosema, 25.  
 catharota, 443.  
 catharurga, 34.  
 cathedraea, 85.  
*Cathelotis*, 241.  
 cathosiota, 200.  
 cato, 374.  
*Catolbistis*, 624.  
 caustocoma, 425.  
*caustogramma*, 366.  
 cedrinopa, 386.  
 celidotis, 169.  
 centracma, 4.  
 centrophracta, 116.  
*Ceranthes*, 317.  
 cerinura, 47.  
 ceriodes, 441.  
*Ceromitia*, 79, 572.  
*Cerostoma*, 87, 360.  
 chalarodes, 153.  
 chalarodesma, 359.  
*Chalceopla*, 53.  
 chalcocrates, 574.  
*Chalcomima*, 507.  
*Chamanthedon*, 267.  
 charadritis, 65.  
 charigramma, 502.  
 charipepla, 326.  
 charixantha, 435.  
*Chelaria*, 30, 282, 513.  
 chelias, 342.  
*Chersomorpha*, 243.  
*Chezala*, 310.  
 chilosema, 219.  
*chionastra*, 313.  
*chionodora*, 192.  
*chionomima*, 488.  
*chionophanes*, 569.  
*chionoptila*, 229.  
*chionotarsa*, 148.  
*chlaniidota*, 355.  
*chlidanopa*, 351.  
 chlorantis, 462.  
*chloraspis*, 106, 369.  
*chlorocrana*, 348.  
*chloroderces*, 487.  
*chlorodeta*, 393.  
*chlorogenes*, 470.  
*chloromis*, 195.  
*chloronephes*, 91.  
*chloroneura*, 6.  
*chlorophthalma*, 32.  
*chloroschema*, 21.  
*chlorosperma*, 528.  
*chlorospora*, 71.  
*chlorostoma*, 478.  
*chloroxantha*, 216.  
*cholerota*, 387.  
*Choreutis*, 305.  
*chorica*, 302.  
*Choronomia*, 317.  
*Choropleca*, 65.  
*Chresmarcha*, 110.  
*chromolitha*, 203.  
*chromotechna*, 212.  
*chrysampyx*, 161.  
 chrysaula, 377.  
*chrysidiás*, 463.  
*chrysobela*, 395.  
*chrysocarpa*, 378.  
*Chrysoclista*, 394.  
*chrysocrana*, 586.  
*chrysodetis*, 248.  
*chrysomelopa*, 369.  
*chrysometra*, 273.  
*chrysomochla*, 90.  
*chrysosoma*, 387.  
*chrysoxesta*, 96.  
*ciliella Busck*, 471.  
*cinnabarina*, 3.  
*cinnabaritis*, 451.  
*circensis*, 459.  
*cirrhocosma*, 627.  
*cirrhocrena*, 517.  
*cirrhocrotala*, 410.  
*cirrholeuca*, 421.  
*cirrhophaea*, 102.  
*citharoeda*, 479.  
*citranches*, 15.  
*citrinopis*, 378.  
*citriseca*, 400.  
*citrochrysa*, 582.  
*citrochyta*, 247.  
*citroclista*, 576.  
*citrocoma*, 72.  
*citroglypta*, 386.  
*citronympha*, 262.  
*citrura*, 371.  
*Cladophantis*, 160, 366.  
*clandestina*, 19.  
*clarisignis*, 93.  
*clastopetra*, 412.  
*claudescens*, 177.  
*cleopatra*, 166.  
*clepsidoxa*, 499.  
*Clepticodes*, 332.  
*climacista*, 289.  
*clistogramma*, 281.  
*clistrodoma*, 21.  
*clitellaria*, 11.  
*clodiana*, 384.  
*clonomicta*, 381.  
*clotho*, 593.  
*Clyisia*, 435.  
*clysmographa*, 197.  
*clytemnestra*, 56.  
*clytozona*, 69.  
*cnemosaris*, 181.  
*Cnephasia*, 116.  
*codonectis*, 339.  
*Coelostathma*, 259.  
*coenulenta*, 355.  
*coercita*, 42.  
*Coleophora*, 66, 374,  
     561, 625.  
*coleoxantha*, 122, 595.

- collega, 364.  
 collegialis, 138.  
 collinearis, 349.  
*Colobocrossa*, 83.  
*Colonanthes*, 12.  
*Colpomorpha*, 528.  
 colposaris, 185.  
 columella, 373.  
 comastis, 316.  
 comissata, 18.  
 comitialis, 138.  
*Commatica*, 503.  
 commixta, 468.  
 commodata, 44.  
 commotica, 61.  
*Commotrias*, 73.  
 commutata, 233.  
*Comocritis*, 118.  
*Comodica*, 82.  
 comparanda, 506.  
 compsocharis, 218.  
*Compsolechia*, 292,  
     353, 506.  
 compsomorpha, 491.  
 compsoneura, 217.  
 compsoptila, 56.  
 compsoxantha, 95.  
 concavata 614.  
 concisa, 505.  
 concitata, 395.  
 congelata, 236.  
*congressella* Walk.,  
     192.  
 congrua, 213.  
*coniographa*, 430.  
 conioleuca, 291.  
 coniopa, 184.  
 conjunctana Mn., 591.  
*conobathra*, 403.  
 constituta, 191.  
 constricta, 226.  
 consummata, 47.  
 contigua, 259.  
 conturbatella Walk.,  
     192.  
 conveniens, 217.  
 convexa, 36.  
 conviciata, 487.  
*Cophomantis*, 290, 356,  
     525.  
*Copromorpha*, 244, 590.  
 coracopis, 324.  
 coronopis, 416.  
*Corsocasis*, 544.  
 corymbitis, 292.  
 corystopa, 382.  
 corythota, 519.  
 cosmangela, 57.  
*Cosmopteryx*, 89, 384,  
     395.  
*Cotyloscia*, 3.  
*Cranaodes*, 327.  
*Crasimorpha*, 33.  
 crassa, 177.  
*Craterobathra*, 379.  
 crateropa, 555.  
 craterospila, 46.  
*crateroxantha*, 321.  
 cratophanes, 522.  
 credula, 343.  
 cremastis, 194.  
*cremastropis*, 599.  
 crescents, 300.  
 cretulata, 345.  
 criminata, 37.  
*Critoxyena*, 623.  
 criopis, 445.  
*crobylistis*, 265.  
 crocacma, 542.  
*Crocanthes*, 519.  
 crococarpa, 385.  
*crocochorda*, 151.  
 crocodeta, 323.  
*crocodilitis*, 554.  
*crocogramma*, 49.  
*crocoleuca*, 51.  
*croconeura*, 267.  
*crocosema*, 524.  
*crocosticta*, 219.  
*Cronicombra*, 306.  
*Crossobela*, 34.  
*crossotorna*, 503.  
*crotalista*, 478.  
 cruda, 311.  
*cryeranthes*, 152.  
*cryeropis*, 167.  
*cryptsetaera*, 183.  
*cryptsigenes*, 523.  
*cryptsrapha*, 581.  
*cryptsinoma*, 527.  
*cryptsiphaea*, 190.  
*Cryptsithyris*, 71, 320,  
     427.  
*cryptsithias*, 558.  
*cryptochlora*, 590.  
*Cryptolechia*, 105, 317,  
     383, 476, 578.  
*Cryptophasa*, 146, 433.  
*cryptosema*, 544.  
*crystallodes*, 539.  
*Ctenocompa*, 328.  
*culminicola* Staud.,  
     323.  
*cultrix*, 87.  
*curialis*, 149.  
*custodita*, 470.  
*Cyanarmostis*, 380.  
*cyanochrysa*, 571.  
*Cyanocrates*, 155.  
*cyanodoxa*, 105.  
*cyanogama*, 584.  
*cyanograptia*, 449.  
*cyanophaea*, 339.  
*cyanoschista*, 289.  
*cyanospira*, 465.  
*cyanospora*, 392.  
*cyanostacta*, 611.  
*cyanozona*, 26.  
*Cyathaula*, 428.  
*cyceonodes*, 278.  
*cycladopa*, 326.  
*cycnolopha*, 186.  
*cynnomorpha*, 169.  
*cyclicota*, 294.  
*cylindrodes*, 84.  
*cymataula*, 320.  
*cymbalistis*, 268.  
*cymogramma*, 193.  
*cymographa*, 536.  
*cymoptila*, 514.  
*Cymotricha*, 1, 287,  
     355, 511.  
*Cynomastix*, 548.  
*Cyphothyris*, 616.  
*cyprantha*, 111.  
*cypodeta*, 556.  
*Cyrictodes*, 283.  
  
 daphnaea, 110.  
 dasyleuca, 242.  
*Decadarchis*, 64, 83,  
     401.  
 decens, 129.  
 dedicata, 133.  
 deferens, 324.  
 defigurata, 407.  
 definita, 438.  
 defixa, 481.  
*deflorescens*, 621.  
*deformata*, 458.  
*defricta*, 253.  
*dehiscens*, 120.  
*delacma*, 24.  
*delatrix*, 17.  
*delia*, 110.  
*delodelta*, 561.  
*delophanta*, 623.  
*delphinodes*, 222.  
*Deltobathra*, 55.  
*deltocausta*, 484.  
*deltochlaena*, 591.  
*deltocrates*, 352.  
*deltographa*, 460.  
*deltomis*, 206.  
*deltoptilus*, 565.  
*deltozela*, 92.  
*deltozyga*, 440.  
 demens, 309, 533.  
*Demobrotis*, 324.

- democratica, 531.  
 demodes, 74.  
 deplanata, 413.  
 Depressaria, 311, 382,  
     475, 621.  
 deridens, 162.  
 Deroxena, 291.  
 descripta, 80.  
 desidiosa, 201.  
 desiliens, 23.  
 detrita, 450.  
 Deuterocopus, 565.  
 devexa, 248.  
 diaconalis, 495.  
 Diactenis, 458.  
 diametrica, 239.  
 dianthes, 439.  
 diasticta, 271.  
 Diceillitis, 460.  
 Dichomeris, 35, 285,  
     354, 510.  
 diclera, 155.  
 dictyophanes, 254.  
 difficilis, 456.  
 diligenda, 477.  
 dilinopa, 198.  
 dimetra, 95.  
 Dinochora, 69.  
 Diocosma, 621.  
 diogma, 336.  
 diophanes, 384.  
 diplanetia, 619.  
 diplospila, 420.  
 diplosticha, 318.  
 diplotima, 269.  
 dirupta, 39.  
 discerpta, 78.  
 disclusa, 16.  
 discrepans, 199.  
 discriminata, 10.  
 disjecta Z., 192.  
 disparata, 24.  
 dispulsa, 309.  
 dissona, 191.  
 distigmatella Z., 525.  
 divulsa, 63.  
 doctissima, 118.  
 Dolerothera, 265.  
 dolichoscia, 394.  
 Dorata, 429.  
 doris, 593.  
 Dorodoca, 90.  
 dorothea, 103.  
 doxanthes, 498.  
 dracuncula, 428.  
 drepanota, 406.  
 Drimylastis, 70.  
 dromica, 185.  
 droserantha, 611.  
 drosocrypta, 273.  
 drosoptila, 82.  
 dryaula, 216.  
 dryocarpa, 142.  
 dryogramma, 567.  
 dryosphaera, 231.  
 ductifera, 329.  
 Durrantia, 161, 364.  
 ebenocista, 433.  
 echinodes, 263.  
 echioglossa, 385.  
 echyropis, 562.  
 eclipticopa, 415.  
 Ectoedemia, 462.  
 effeta, 117.  
 ejiciens, 412.  
 Elachista, 60, 296.  
 elaeanthes, 340.  
 elaeurga, 226.  
 elaphodes, 618.  
 elaphopa, 294.  
 Elaphrerga, 574.  
 electrantha, 377.  
 elimata, 611.  
 embolarcha, 500.  
 embrithopa, 366.  
 emicans, 18.  
 emollita, 234.  
 empedarthra, 124.  
 Empedaula, 11.  
 empedocles, 552.  
 Emphantica, 430.  
 encamina, 379.  
 encomias, 376.  
 Encrasima, 50.  
 endobela, 296.  
 endocarpa, 419.  
 endochra, 198.  
 endoclina, 426.  
 endoneurias, 134.  
 endophaea, 567.  
 endopyrrha, 609.  
 Endothetis, 549.  
 endoxa, 263.  
 enervata, 532.  
 enoptrias, 508.  
 enormis, 394.  
 ensigera, 150.  
 enstacta, 429.  
 Enthetica, 38.  
 enthrypta, 397.  
 Eochrois, 574.  
 Eomichla, 469.  
 eoplecta, 142.  
 Epactris, 72.  
 Epermenia, 52, 626.  
 Epicephala, 407, 579,  
     583.  
 epichares, 506.  
 Epicharta, 285.  
 epiclentes, 452.  
 epicremna, 251.  
 Epidola, 270.  
 epigypsa, 351.  
 Epimecyntis, 100, 470.  
 Epimoryctis, 618.  
 epiphanta, 284.  
 Epiphractis, 577.  
 episcopopa, 268.  
 Episimus, 337.  
 Epithestis, 15, 275,  
     486.  
 eporphna, 62.  
 erasa, 442.  
 erebocrossa, 574.  
 erebodoxa, 351.  
 Eretmocera, 380, 543,  
     586.  
 eridarcha, 336.  
 eridopa, 407.  
 Eriogenes, 159.  
 erotopis, 315.  
 ersodes, 367.  
 erythranches, 269.  
 erythrites, 461.  
 erythrocyma, 612.  
 erythrodesma, 595.  
 erythroleuca, 473.  
 erythromis, 111.  
 eschatopa, 100.  
 essedaria, 306.  
 eteoxyantha, 533.  
 Ethmia, 119, 362, 418,  
     563.  
 euarithma, 120.  
 eualypta, 436.  
 eucausta, 492.  
 eucentrota, 574.  
 euchorda, 31.  
 eucolapta, 73.  
 eucoma, 168.  
 Eucosma, 67, 140, 333,  
     439, 599.  
 euglypta, 368.  
 Eulia, 249, 610.  
 eulidias, 246.  
 eumarodes, 68.  
 Eumenodora, 91.  
 eumeris, 9.  
 eumetrota, 319.  
 Eumiturga, 177, 233.  
 Eunebristis, 3.  
 Eunomarcha, 26.  
 euophthalma, 83.  
 Eupetochira, 363.  
 euphanopis, 377.  
 euphrantis, 325.  
 euphonopa, 370.  
 Euplocamus, 74.  
 euprosopa, 276.

*eurychlora*, 258.  
*eurycitra*, 463.  
*eurycrates*, 121.  
*eurycryptis*, 311.  
*eurydryas*, 314.  
*eurylyca*, 417.  
*eurymenes*, 608.  
*eurymolybda*, 306.  
*eurynephela*, 534.  
*euryperena*, 597.  
*euryplintha*, 53.  
*euthalama*, 112.  
*euthetodes*, 509.  
*Euthorybeta*, 132.  
*euthyrsa*, 577.  
*Euxanthis*, 436, 591.  
*Evetria*, 333.  
*evocata*, 53.  
*exaequata*, 52.  
*exaeta*, 264.  
*exagitata*, 319.  
*exallacta*, 287.  
*exanthistis*, 619.  
*excors*, 568.  
*excreta*, 568.  
*exempta*, 179.  
*exilis Wals.*, 343.  
*exorycha*, 409.  
*expleta*, 54.  
*explicata*, 510.  
*exsecta*, 354.  
*extans*, 272.  
*exstigmata*, 468.  
*externella Walk.*, 192.  
*extorris*, 50.  
  
*fabivora*, 449.  
*fabricata*, 242.  
*faeculosa*, 453.  
*fasciformis*, 582.  
*fercularia*, 492.  
*ferruginea*, 137.  
*festicola*, 104.  
*Filinota*, 312, 574.  
*finita*, 116.  
*finitrix*, 208.  
*fissirostris*, 413.  
*fistularis*, 449.  
*flaccescens*, 161.  
*flagellatrix*, 499.  
*flaminia*, 255.  
*flaviceps Feld.*, 423.  
*flavidorsis*, 573.  
*flocculosa*, 178.  
*florescens*, 141.  
*floricoma*, 123.  
*fluctuans*, 2.  
*foederalis*, 43.  
*formularis*, 503.

*Fortinea*, 270.  
*Frisilia*, 517.  
*frustulenta*, 13.  
*fuliginea*, 490.  
*fuliginella Dup.*, 583.  
*fulta*, 234.  
*funicularis*, 228.  
*furcigera*, 460.  
*furculata*, 101.  
*fustigera*, 408.  
  
*Gaes*, 286, 510.  
*galactacma*, 89, 117.  
*galactarcha*, 418.  
*galatea*, 278.  
*Galtica*, 270.  
*Gelechia*, 19, 276, 345,  
 488.  
*Gemorodes*, 154.  
*generatrix*, 239.  
*geniculata*, 591.  
*genitrix*, 380.  
*gennaea*, 62.  
*geodes*, 515.  
*geraropa*, 371.  
*germinans*, 211.  
*Gerontha*, 428.  
*glyceranthes*, 445.  
*glycinopis*, 26.  
*glycitis*, 438.  
*Glycynympha*, 155.  
*glycystrota*, 462.  
*Glyphidocera*, 48, 529.  
*glyphidopa*, 475.  
*Glyphipteryx*, 136,  
 308, 424.  
*glyptocosma*, 99.  
*gnomonodes*, 285.  
*Gnorimoschema*, 491.  
*Gnosimacha*, 354.  
*gomphacma*, 440.  
*Gomphocrates*, 288.  
*gomphopis*, 350.  
*Gonada*, 98, 593.  
*goniometra*, 522.  
*Gracilaria*, 86, 295, 409  
 480, 582.  
*gracula*, 495.  
*grammotorna*, 256.  
*grammozona*, 135.  
*grata*, 472.  
*gravescens*, 237.  
*griseicoma*, 115.  
*gubernatrix*, 173.  
*Gymnelema*, 78.  
*Gymnogelastis*, 587.  
*Gymnogramma*, 120,  
 597.

*gymnolopha*, 174.  
*gypsopsis*, 474.  
  
*haemanthes*, 370.  
*haematantha*, 121.  
*haemodryas*, 546.  
*haemogastra*, 297.  
*haemograptis*, 451.  
*Haemolytis*, 245.  
*haemorrhanta*, 105.  
*haemorrhoa*, 383.  
*haemothicta*, 257.  
*halans*, 92.  
*haligramma*, 282.  
*haliplancta*, 325.  
*halitheia*, 330.  
*halmas*, 200.  
*halmyropis*, 270.  
*halobapta*, 575.  
*halycopa*, 350.  
*hamaxastra*, 563.  
*hapalosarca*, 67.  
*haplocentra*, 170.  
*Haplochela*, 32.  
*haplodoxa*, 162.  
*Hapsifera*, 553.  
*hebaea*, 89.  
*Hebdomactis*, 539.  
*Hecatopeda*, 543.  
*Hectacma*, 84.  
*Hedroxena*, 92.  
*Helcystogramma*, 26.  
*helena*, 388.  
*heliaspis*, 303.  
*helicomitra*, 85.  
*helicosema*, 240.  
*helicosticta*, 490.  
*heliochalca*, 464.  
*heliochrosta*, 437.  
*heliographa*, 520.  
*heliomima*, 576.  
*heliotoma*, 267.  
*helophaea*, 442.  
*hemeropa*, 2.  
*hemiclina*, 512.  
*hemicrates*, 613.  
*hemicrossa*, 345.  
*hemicryptis*, 414.  
*Hemimene*, 341.  
*hemiphanta*, 198.  
*hemiphracta*, 320.  
*hemispora*, 81.  
*hemistacta*, 85.  
*heptanoma*, 138.  
*heptaspila*, 589.  
*hercoptila*, 333.  
*heroica*, 391.  
*Heteralcis*, 517.  
*heteranthes*, 296.

- heteraula, 360.  
 Heteroderces, 521.  
 heterodesma, 547.  
 heterograpta, 425.  
 Heterogymna, 138.  
 Heterotactis, 392.  
 hexacopa, 483.  
 hexanesa, 501.  
 hexascia, 215.  
 Hexeretmis, 539.  
 Hierangela, 498.  
 hieraspis, 89.  
 hierograpta, 139.  
 Hieromantis, 66, 541.  
 Hierophanes, 586.  
 hierophanes, 153.  
 Hieroxestis, 63.  
 Hilarographa, 627.  
 hippophylax, 572.  
 hipposaris, 348.  
 hippotarcha, 41.  
 holarga, 160.  
 Holcopogon, 515.  
 holocitra, 156.  
 holoperca, 78.  
 holotephras, 67.  
 holotorna, 157.  
 homalocrossa, 586.  
 Homalopsycha, 70.  
 Homaloxestis, 518.  
 Homona, 106, 455, 605.  
 homosacta, 610.  
 hoplanthes, 75.  
 hoplista, 340.  
 hoplitica, 216.  
 hoplodoxa, 507.  
 horizontias, 188.  
 Hormantris, 327.  
 hormista, 409.  
 hormocrossa, 148.  
 horocentra, 332.  
 horochlora, 519.  
 horocyma, 209.  
 horographa, 454.  
 horometra, 222.  
 horoscopa, 279.  
 hyacinthitis, 559.  
 Hyalopseustis, 157.  
 hyaloxantha, 467.  
 Hydarantes, 460.  
 hydrochoa, 610.  
 hydronoma, 561.  
 hydrophora, 171.  
 hygraema, 361.  
 hylarcha, 400.  
 hylophanes, 447.  
 hyperbolica, 471.  
 Hypercallia, 103, 312,  
     475, 575, 621.  
 Hyperecta, 508.  
 Hypersymmoca, 63.  
 Hypochalcia, 583.  
 Hypochasmia, 517.  
 hypochrysa, 332.  
 Hyponomeuta, 117,  
     415.  
 Hypophrictis, 548.  
 hypsicrates, 130.  
 hypsidryas, 140.  
 hypsipola, 290.  
 hypsisthrona, 251.  
 ichthyodes, 231.  
 icteropis, 209.  
 idalia, 62.  
 Idiophantis, 24.  
 Idioptila, 343.  
 igniflora, 606.  
 Ilingiotis, 2.  
 illicita, 511.  
 illita, 232.  
 illudens, 76.  
 illuminatrix, 537.  
 imbrifera, 538.  
 Imma, 132, 302, 375,  
     420.  
 immutabilis, 259.  
 imparata, 430.  
 improvida, 275.  
 inardescens, 202.  
 incantata, 469.  
 incitatrix, 207.  
 inclemens, 466.  
 inclusa, 355.  
 inclyta, 144.  
 inconcinna, 59.  
 incrustata, 553.  
 indefessa, 528.  
 indentatus, 564.  
 indicatrix, 84.  
 indiserta, 285.  
 inerudita, 290.  
 infamis, 60.  
 infervescens, 596.  
 infusoria, 259.  
 injucunda, 219.  
 innexa, 190.  
 inophora, 428.  
 inota, 87.  
 insecuratrix, 127.  
 insimulata, 237.  
 insipiens, 11.  
 instabilis, 75.  
 insulata, 115.  
 intercisus, 565.  
 interfusa, 330.  
 interjecta, 54.  
 interscissa, 76.  
 introspiciens, 287.  
 intuens, 59.  
 inurbana, 49.  
 inventrix, 156.  
 inviolata, 152.  
 iobapta, 352.  
 iocapna, 154.  
 ioplantha, 600.  
 iospora, 482.  
 iostalacta, 212.  
 ioxanthas, 599.  
 iphicleia, 98.  
 iras, 237.  
 Irianassa, 301.  
 iricolor, 601.  
 iridesma, 598.  
 irrisoria, 76.  
 irrorella Woll., 321.  
 isarga, 224.  
 isastræ, 312.  
 ischioptila, 211.  
 ischnodesma, 464, 625.  
 Ischnodoris, 528.  
 ischnoscia, 401.  
 Isembola, 271.  
 isocharis, 547.  
 isoclista, 136.  
 isocrates, 94.  
 Isocrita, 382.  
 isocrossa, 346.  
 Isocylindra, 584.  
 isodoxa, 444.  
 isographa, 176.  
 isograpta, 410.  
 isomacha, 132.  
 isonephes, 531.  
 isoniphæs, 88.  
 isopela, 153.  
 isopelta, 482.  
 isopercna, 340.  
 isophrictis, 481.  
 isoplasta, 516.  
 isoplantha, 193.  
 isopoca, 551.  
 isopogon, 513.  
 Isorrhœa, 65, 96.  
 isorrhithma, 262.  
 isostacta, 273.  
 isothea, 615.  
 isotherma, 617.  
 isoxysta, 317.  
 ithyclina, 260.  
 ithycypha, 373.  
 ithymetra, 312.  
 itygramma, 422.  
 jactatrix, 272.  
 jason, 568.  
 javarica Butl., 192.  
 juncta, 258.  
 juvenilis, 481.

- Knabi Wals., 547.  
 Labdia, 56, 90, 384, 545.  
 laconica, 365.  
 Lactura, 121, 415, 594.  
 lagaropis, 474.  
 lagodes, 278.  
 lamprocrossa, 400.  
 lampropoda Hamps., 585.  
 Lamproteucha, 585.  
 lasiandra, 143.  
 Laspeyresia, 68, 144, 342, 447, 604.  
 lathiptila, 190.  
 laticivora, 372.  
 lavata, 550.  
 lechriosema, 475.  
 Lecithocera, 38, 290, 522.  
 lecticata, 282.  
 lectulifera, 493.  
 Leistarcha, 574.  
 Leistogenes, 363.  
 leptaula, 360.  
 leptobrocha, 44.  
 leptochorda, 299.  
 leptoclista, 541.  
 leptocrossa, 273.  
 leptodesma, 480.  
 leptogma, 217.  
 leptogrpta, 115, 369.  
 leptophanes, 411.  
 Leptozestis, 91.  
 Lepyrotica, 326.  
 leucactis, 158.  
 leucana Sepp, 192.  
 leucatma, 144.  
 leucobactra, 256.  
 leucobathra, 61.  
 leucocapna, 227.  
 leucodelta, 160.  
 leucodora, 447.  
 Leucogonia, 504.  
 leucograpta, 45.  
 leucomias, 162.  
 leucoplasta, 232.  
 leucopogon, 83.  
 leucoporpa, 313.  
 Leucoptera, 265, 396.  
 leucorectis, 223.  
 leucoritis, 381.  
 leucosaris, 196.  
 leucoscia, 359.  
 leucosticta, 391.  
 leucozostra, 5.  
 Leuronoma, 343.  
 liaropa, 71.  
 liberta, 291.  
 libidinosa, 311.  
 lichenista, 312.  
 lignaria, 292.  
 liliacea, 622.  
 lilitha, 587.  
 Limenarchis, 288.  
 limenias, 437.  
 Limnoecia, 59, 90, 392, 546.  
 linoplecta, 67.  
 linozona, 269.  
 liobathra, 55.  
 liophaea, 361.  
 Liopseustis, 429.  
 liparochra, 452.  
 liquescens, 227.  
 lithochalca, 581.  
 lithochrysa, 580.  
 Lithocolletis, 406, 582,  
 lithodoxa, 529.  
 Lobesia, 442.  
 Locharcha, 18.  
 locharcha, 94.  
 Logisis, 270.  
 longa, 381.  
 longifasciella Clem., 345.  
 longinqua, 25.  
 lophandra, 530.  
 Lophoprora, 611.  
 lophoptycha, 188.  
 lophosaris, 186.  
 loranthivora, 591.  
 loxaula, 396.  
 loxochorda, 315.  
 lucrosa, 184.  
 lunularis, 310.  
 luticoma, 489.  
 lutraula, 20.  
 lychnacma, 378.  
 lychnitis, 413.  
 Lychnocrates, 226.  
 lycophanes, 78.  
 Lyonetia, 81.  
 lyroda, 357.  
 Lysipatha, 289.  
 Machimia, 311, 470, 618.  
 Machimostola, 399.  
 machinata, 488.  
 machinatrix, 208.  
 Machlotica, 306.  
 macranthes, 480.  
 macraulax, 559.  
 Macrobathra, 98, 310, 380, 614.  
 macroctenis, 243.  
 macrornis, 567.  
 macrorrhyncha, 592.  
 maenas, 109.  
 magdalena, 622.  
 magnetica, 451.  
 magocosma, 594.  
 maleficus, 296.  
 malevolens, 433.  
 malivora, 555.  
 mallodeta, 73.  
 malthaca, 401.  
 manceps, 172.  
 mansueta, 54.  
 marantica Wals., 129.  
 margaritacea, 104.  
 Marmara, 480.  
 marmarantha, 109.  
 marmarozona, 421.  
 maschalista, 292.  
 Mastigostoma, 330.  
 maturescens, 223.  
 meditans, 504.  
 megacentra, 59.  
 Megacraspedus, 272, 343.  
 megaleuca, 224.  
 megalochra, 570.  
 melanaphra, 10.  
 melanaspis, 337.  
 melanchlora, 496.  
 melanchropa, 281.  
 melanixa, 192.  
 melanocarpa, 527.  
 melanoclista, 362.  
 melanoloncha, 301.  
 melanometra, 229.  
 melanoplantha, 279.  
 melanortha, 510.  
 melanothicta, 367.  
 Melanozestis, 547.  
 Melasina, 76, 328, 430, 550.  
 Meleonoma, 623.  
 meligrapta, 183.  
 melinopa, 222.  
 meliphaga, 64.  
 meliptila, 283.  
 meliscelis, 542.  
 Melisophista, 371.  
 melitocrossa, 253.  
 Melitonympha, 360.  
 melitoploca, 329.  
 melitoptila, 17.  
 melitopyga, 41.  
 melitosoma, 584.  
 Melittia, 467.  
 melixesta, 210.  
 melobaphes Wals., 314.  
 Melodryas, 131, 593.  
 meloxantha, 364.

- mentigera, 236.  
*Meridarchis*, 137, 404,  
  588.  
*merocentra*, 147.  
*mesatma*, 269.  
*mesobactris*, 590.  
*mesocrossa*, 379.  
*mesogypsa*, 159.  
*mesolampra*, 376.  
*mesosaris*, 178.  
*mesotoma*, 149.  
*mesotricha*, 337.  
*Metabolaea*, 32.  
*metacapna*, 108.  
*metadesma*, 352.  
*Metallocrates*, 579.  
*metamorphica*, 439.  
*metaphracta*, 156.  
*metaspila*, 302.  
*Meteoristis*, 27.  
*metrionoma*, 470.  
*micrarcha*, 261.  
*microcasis*, 532.  
*Microcolona*, 391.  
*microdesma*, 534.  
*microdictyas*, 58.  
*microglypta*, 393.  
*Microschismus*, 373.  
*milictis*, 163.  
*minerva*, 477.  
*miniana*, 246.  
*minna Busck*, 192.  
*miranda*, 587.  
*miseranda*, 62.  
*mixogama*, 474.  
*mnesicosma*, 119.  
*mnesileuca*, 397.  
*Mnesistega*, 36, 356.  
*mniadora*, 204.  
*mochlopis*, 1.  
*molinda*, 145.  
*mollifluia*, 373.  
*Molopostola*, 284.  
*molybdaspis*, 316.  
*molybdochtha*, 603.  
*Mometa*, 351.  
*Mompha*, 547.  
*monactis*, 10.  
*monelpis*, 398.  
*monoglossa*, 80.  
*monopa*, 350.  
*monophaea*, 380.  
*Monopis*, 71, 321, 425.  
*montezuma*, 323.  
*montivola*, 364.  
*mundata*, 495.  
*munerata*, 66.  
*murenula*, 65.  
*murmurans*, 382.  
*muscicolor*, 560.  
*muscosa*, 335.  
*muscosana*, Zell., 249.  
*musicosema*, 305.  
*Musurga*, 3.  
*Myconita*, 27.  
*myelopis*, 322.  
*mylias*, 420.  
*myloclasta*, 321.  
*myriocosma*, 608.  
*myriodesma*, 533.  
*myrmecias*, 590.  
*Myrmecozela*, 425.  
*myrodora*, 197.  
*myroxesta*, 114.  
*Mysaromima*, 227.  
*mystica*, 563.  
*mysticopis*, 233.  
  
*Narycia*, 328, 430, 554.  
*nauarches*, 569.  
*nausicaa*, 288.  
*Nealyda*, 5.  
*neatodes*, 35.  
*negotiosa*, 201.  
*nematodes*, 452.  
*Nemotois*, 52, 79, 462,  
  571, 624.  
*Neocorodes*, 36.  
*neocrates*, 146.  
*neocrossa*, 193.  
*Neodactylota*, 488.  
*neolopha*, 259.  
*Neophylarcha*, 240.  
*Neopseustis*, 404.  
*neoptila*, 195.  
*neospila*, 471.  
*nephanthes*, 497.  
*nephelaspis*, 276.  
*nephelatma*, 375.  
*nephelodesma*, 280.  
*nephelomicta*, 555.  
*nepheloscopa*, 426.  
*nephelozyga*, 419.  
*nephodryas*, 420.  
*nephospora*, 623.  
*Nepticula*, 89, 117, 295,  
  461.  
*nereidopa*, 333.  
*nesocharis*, 427.  
*nesoclera*, 499.  
*nesographa*, 244.  
*nesograpta*, 150.  
*Nesophylax*, 320.  
*neurocentra*, 179.  
*neurocoma*, 614.  
*neurophanes*, 271.  
*neurotypa*, 586.  
*Nicanthes*, 395.  
*nictitans* Zell., 192.  
  
*oblitescens*, 71.  
*obscura* Butl., 123.  
*obstans*, 413.  
*obtrectans*, 551.  
*obtruncata*, 30.  
*obumbrata*, 156.  
*ochraspis*, 535.  
*ochricollis* Zell., 192.  
*ochrocapna*, 40.  
*ochrochorda*, 118.  
*ochrodeta*, 23.  
*ochromeris*, 398.  
*ochromorpha*, 145.  
*ochrophthalma*, 344.  
*ochrosaris*, 199.  
*ochroschista*, 494.  
*ochrospora*, 472.  
*ochrothicta*, 180.

- ochrotoma, 17.  
 ochroxysta, 483.  
 ochthotoma, 616.  
 Octasphales, 620.  
 octobola, 137.  
 octophora, 508.  
*Octopunctata* Turn.,  
     86, 409.  
 octosticta, 610.  
 odiosa, 93.  
*Odites*, 156, 364, 434,  
     555.  
*Oedematopoda*, 542.  
*Oegoconia*, 291.  
 oenochares, 100.  
 Oenoe, 325.  
 ogmologpha, 557.  
*Oinophila*, 398, 401.  
 oligastraa, 308.  
 omichlopis, 427.  
 ommatopa, 230.  
*Onebala*, 42, 508.  
 ophiodora, 317.  
 ophioglossa, 122.  
 ophitis Wals., 503.  
 ophiucha, 123.  
 ophryodes, 368, 616.  
*Opogona*, 398, 400.  
 oporista, 310.  
*Opotega*, 396.  
 opportuna, 38.  
 oppressa, 258.  
 opsorrhao, 512.  
 opticodes, 370.  
*orbicularis* Feld., 592.  
*Ordrupia*, 242.  
 orgadopa, 182.  
 orgias, 607.  
 orgilopis, 16.  
 orientalis Stt., 579.  
*Orinympha*, 360.  
 orinympha, 304.  
 oriphanes, 608.  
 oriphanta, 473.  
*Orneodes*, 95, 373, 533.  
 orneopis, 221.  
 oroglypta, 595.  
*Orsodytis*, 286.  
 orthiasta, 427.  
*Orthiostola*, 357, 478.  
 Orthochtha, 399.  
 orthoctenis, 49.  
 orthographa, 205.  
 orthomeris, 528.  
 orthomorpha, 443.  
 orthrarcha, 578.  
 orthriopa, 166.  
 orthritis, 545.  
*Orygocera*, 383, 577,  
     622.  
 osphrantica, 525.  
*osseella* Stt., 486.  
 osteosema, 486.  
 ostiaria, 322.  
 ovulifera, 182.  
 oxychasta, 487.  
 oxycryptis, 516.  
 oxylitha, 521.  
 oxyloba, 492.  
 oxyloga, 408.  
*Oxymachaeris*, 331.  
 oxymeris, 486.  
 oxyptila, 389.  
*Oxyptilus*, 93, 296,  
     564.  
*Oxysactis*, 35.  
 oxyschista, 187.  
 oxythectis, 481.  
 oxytricha, 424.  
 oxytypa, 31.  
  
 Pachnistis, 45, 526.  
 pachnostola, 598.  
*Pachygeneia*, 11.  
*Pachyrhabda*, 377.  
 pactota, 164.  
 paeta, 522.  
 palaearcha, 582.  
 palaeocosma, 303.  
 palaeographa, 432.  
 palamota, 274.  
*Palinorsa*, 99.  
 palpalis Zell., 239.  
*Pammene*, 447.  
 panarista Turn., 619.  
 panchromatica, 274.  
 pancyclia, 434.  
 pansarista, 526.  
*Pansepta*, 153.  
*Parachanda*, 338.  
*Paracharactis*, 470.  
 paradelta, 141.  
*Paralecta*, 153.  
 paraleuca, 482.  
 paralonoma, 427.  
*Paranthrene*, 267, 584.  
 paraplecta, 214.  
*Paraptila*, 259, 610.  
 parasitica, 626.  
 parastropha, 454.  
*Paratheta*, 467.  
 parazela, 505.  
*Parectopa*, 409.  
 pariodelta, 541.  
 paroditis, 390.  
 parorthota, 453.  
 patarea, 110.  
 patriocosma, 341.  
 paulina, 140.  
  
 pauperatella Walk.,  
     192.  
*Peleopoda*, 161.  
 pelinactis, 133.  
 pelocrossa, 396.  
 pelodoxa, 431.  
 pelospora, 356.  
 pelostrota, 328.  
 penetrans, 13.  
 pentadisca, 71.  
 pentagama, 388.  
 pentaxantha, 543.  
 peperistis, 277.  
 peplotoma, 442.  
 peragrata, 33.  
 percnocarpa, 181.  
 percnocoma, 578.  
 percnogona, 170.  
 percnoleuca, 48.  
 percnophanes, 543.  
 percnorma, 618.  
*Periacma*, 62, 623.  
 pericharacta, 70.  
*Peridaedala*, 139.  
 peridesma, 220.  
 peridoxa, 97.  
 perilithas, 331.  
 perioditis, 471.  
 perischias, 129.  
 periscias, 434.  
*Peritorneuta*, 62.  
 perjura, 207.  
 perlucescens, 328.  
 permutata, 457.  
*Peronea*, 56, 612.  
 persicopa, 416.  
*Persicoptila*, 60, 390.  
 perstricta, 453.  
 perterrita, 11.  
 pessocrossa, 273.  
*Pessograptis*, 29.  
 petalocrossa, 250.  
*Petasanthes*, 158.  
*Pezetaera*, 74.  
 phaeochtha, 147.  
 phaeodora, 541.  
 phaeomystis, 197.  
 phaeonephela, 322.  
 phaeopleura, 90.  
 phaeosoma, 396.  
 phaeostrota, 35.  
 phaeothicta, 89.  
 phaeoxesta, 111.  
 phalarista, 615.  
*Phalarotarsa*, 101.  
*Phalonia*, 52, 139, 366,  
     438, 591.  
 phanerarcha, 96.  
*Phanerochersa*, 241.  
 phanerodoxa, 291.

- phantastis, 338.  
 Phasmatographa, 418.  
 Philagraula, 554.  
 Philametris, 102.  
 Philausteria, 358.  
 Philedone, 247.  
 philocremna, 570.  
 philomela, 224.  
 philomusa, 527.  
 philonoma, 132.  
 philorectis, 298.  
 phlyctaenopa, 241.  
 phoenocrossa, 141.  
 phoenoteles, 519.  
 phenoxyantha, 60.  
 phormophora, 284.  
 photodoxa, 571.  
 phratriastis, 523.  
 phrixopa, 264.  
 phronimopis, 390.  
 Phruriastis, 64.  
 phryactis, 167.  
 Phtheochroa, 367.  
 Phthorimaea, 23, 50,  
     278, 493, 532.  
 phthoromima, 617.  
 Phyllocnistis, 262, 405,  
     579.  
 phyllograptia, 476, 622.  
 Phylopatris, 14.  
 physocoma, 286.  
 picrocosma, 265.  
 picrogramma, 489.  
 picromorpha, 469.  
 picroscopa, 588.  
 Picrotechna, 317.  
 Piloprepes, 617.  
 pindarica, 118.  
 pinguicula, 489.  
 piratica, 414.  
 piratodes, 601.  
 Pisistrata, 81.  
 Placanthes, 42.  
 planicola, 429.  
 planicomæ, 189.  
 platamodes, 55.  
 platyantha, 113.  
 platycosma, 582.  
 platycypha, 418.  
 platydoxa, 20.  
 Platydæra, 351.  
 platyloxa, 553.  
 platymolybdis, 603.  
 platyorma, 122.  
 Platytilla, 93, 297,  
     566.  
 platyrrhoa, 61.  
 platyrrhyncha, 402.  
 plebigena, 78.  
 plectanopa, 12.  
 plerotis, 165.  
 pleuracma, 280.  
 pleurosaris, 14.  
 Pleurota, 61, 473.  
 pleurotricha, 206.  
 poecilantha, 113.  
 pogonias, 43.  
 politica, 532.  
 polyaema, 4.  
 polyantha, 69.  
 polyarcha, 106.  
 Polychrosis, 142, 443.  
 Polydrachma, 461.  
 polygrapta, 82.  
 Polyhymno, 280, 502.  
 Polymnestra, 331.  
 polytechna, 114.  
 pompeiana, 176.  
 Pompostola, 325.  
 pontifraga, 444.  
 porphyraspis, 625.  
 porphyropa, 332.  
 porphyrospila, 306.  
 potamites Wals., 108.  
 praedicata, 19.  
 praeducta, 511.  
 praenivalis, 152.  
 praenubila, 231.  
 praesiliens, 68.  
 praestabilis, 323.  
 praetexta, 79.  
 prasinaula, 359.  
 Prasodryas, 287.  
 pratifera, 222.  
 Prays, 361, 419.  
 primipilana, 8.  
 prionistis, 457.  
 prionota, 407.  
 prismatica Wals., 324.  
 pristina, 160.  
 privata, 121.  
 Proadamas, 527.  
 Problastodes, 399.  
 Probolacma, 362.  
 probolaspis, 508.  
 probolopis, 8.  
 proceros, 422.  
 procritica, 217.  
 Prodidactis, 563.  
 projectrix, 549.  
 prolectans, 314.  
 Promalactis, 61, 615.  
 Promenesta, 161.  
 Promolopica, 284.  
 promonitrix, 345.  
 Pronomeuta, 562.  
 propycnota, 577.  
 prorepta, 19.  
 prosarithma, 425.  
 Proschistis, 601.  
 prosiliens, 454.  
 prostylas, 327.  
 Protanystis, 97.  
 protaxia, 72.  
 Prothinodes, 70.  
 protophanes, 382.  
 Prototheora, 80.  
 protracta, 74.  
 prudentula, 232.  
 prymnaea, 619.  
 psacasta, 483.  
 psaliodes, 294.  
 psaltria, 479.  
 Psamathocrita, 486.  
 psammochtha, 147.  
 psammocryptis, 63.  
 psammolitha, 48.  
 Psegmatica, 598.  
 psephophragma, 518.  
 Pseudatteria, 108, 606.  
 Pseudodoxia, 62, 478.  
 Pseudurgis, 74.  
 psilodoxa, 28.  
 Psittacastis, 316.  
 Psychromnestra, 88.  
 psyllodecta, 121.  
 Pterolonche, 87.  
 Pterophorus, 299, 568.  
 pterygota, 503.  
 Ptilogenes, 228, 433.  
 Ptilothyris, 288.  
 ptilozona, 64.  
 ptochodyras, 46.  
 ptochomorpha, 36.  
 Ptochoryctis, 152.  
 pugnax, 266.  
 pulcherrima Walk.,  
     619.  
 pulvinata, 128.  
 pumicata, 525.  
 punica, 554.  
 purpurissatana Kenn.,  
     604.  
 putris, 60.  
 pycnochra, 448.  
 pycnostrota, 600.  
 Pyramidobela, 414.  
 pyrangela, 400.  
 pyrocanthara, 112.  
 pyrocephala 543.  
 Pyroderces, 59, 90,  
     389, 545.  
 pyrodoxa, 310.  
 pyrogramma, 84.  
 pyrometalla, 401.  
 pyromyia, 543.  
 pyronota, 98.  
 pyronympha, 127.  
 pyrophracta, 97.  
 Pyrophractis, 617.

- pyrostoma, 372, 465.  
 pyrothorax, 415.  
 pyrrhocolona, 257.  
 pyrrhogamma, 585.  
  
 quadratella Walk., 192  
 quadrilineella Chamb.,  
     384.  
 quincuncialis, 392.  
 quinquepunctella  
     Wals., 507.  
  
 rabida, 500.  
 ramulata, 278.  
 rebellis, 363.  
 reciprocans, 95.  
 reclusa, 53.  
 recrepans, 310.  
 rectificata, 180.  
 recurrens, 210.  
 Recurvaria, 13, 484.  
 recurvata, 39.  
 recurvella Walk., 192.  
 redintegrata, 218.  
 reflectrix, 448.  
 refrigerescens, 68.  
 regesta, 225.  
 relapsa, 444.  
 relata, 207.  
 religiosa, 28.  
 reliquatrix, 436.  
 remorsa, 195.  
 renascens Wals., 510.  
 reprehensa, 238.  
 repudiata, 6.  
 residua, 41.  
 residuella Zell., 558.  
 resignata, 510.  
 resiliens, 173.  
 rhabdophanes, 396.  
 Rhapsodica, 363.  
 rhedaria, 20.  
 rhipidaspis, 390.  
 rhipidoma, 142.  
 rhizonympha, 90.  
 rhodanopa, 358.  
 rhodochropa, 369.  
 rhodoclea, 395.  
 rhodoglaucia, 620.  
 rhodographa, 417.  
 rhodopola, 621.  
 rhombophora, 265.  
 Rhopalosetia, 241.  
 rhothodoxa, 548.  
 Rhynchotona, 35.  
 rhythmodes, 506.  
 rhythmogramma, 106.  
 Rhythmologa, 249.  
  
 rhythmopis, 431.  
 rivalis, 281.  
 rivulata, 348.  
 rixata, 80.  
 robiginosa, 215.  
 rubriflora, 596.  
 ruens, 538.  
  
 sacrificus, 299.  
 saeva, 566.  
 Sagalassa, 132.  
 sagittans, 33.  
 saliaris, 459.  
 saliens, 385.  
 salinatrix, 273.  
 salome Busck, 192.  
 salubris, 208.  
 sana, 235.  
 sandycopa, 155.  
 sanguiflua, 123.  
 sanguipennis, 267.  
 sanidopa, 242.  
 sarcochlora, 491.  
 sarcocrossa, 86.  
 sarcopis, 562.  
 sarcostega, 107.  
 sardania, 168.  
 Sarisophora, 36.  
 satelles, 202.  
 Sathrogenes, 2.  
 satyrisca, 329.  
 saxeae, 14.  
 scaeodesma, 409.  
 Scaeosopha, 469.  
 scandalota, 294.  
 scardamycitis, 326.  
 sceliphrodes, 150.  
 scenodoxa, 57.  
 sceptrophora, 276.  
 Schoenotenes, 116, 459,  
     610.  
 scholasticus, 94.  
 sciactis, 38.  
 sciaphilina Zell., 192.  
 Scieropepla, 154.  
 sciochalca, 413.  
 sciocnesta, 213.  
 sciodelta, 578.  
 sciodyras, 255.  
 sciogama, 558.  
 sciomochla, 430.  
 scioneura, 515.  
 sciota, 353.  
 scitiarella Walk., 192.  
 scitissimella Walk.,  
     574.  
 Sclerograptis, 31.  
 scoleciias, 419.  
 Scoparia, 583.  
  
 Scorpionsis, 619.  
 scribarius, 300.  
 scutellata, 230.  
 scutifer, 564.  
 Scythriss, 87, 412, 561.  
 scythropis, 583.  
 Secitis, 472.  
 seclusa, 247.  
 secundata, 214.  
 sejuncta, 517.  
 selastis, 261.  
 selenaspis, 133.  
 sellifera, 163.  
 semifervens, 379.  
 semiovata, 234.  
 semiramis, 545.  
 semirosea, 492.  
 semisecta, 393.  
 semitincta, 562.  
 semnias, 286.  
 semnocrana, 433.  
 semnolitha, 386.  
 semophantis, 537.  
 sensilis, 467.  
 sepulcrana, 334.  
 serarcha, 556.  
 Sericostola, 358.  
 serriformis, 580.  
 sertigera, 27.  
 sidiota, 96.  
 signigera, 358.  
 silvana, 560.  
 Simaethis, 304.  
 Simoneura, 503.  
 Sirogenes, 3.  
 Sitotroga, 483.  
 smaragdaspis, 79.  
 smaragdopis, 275.  
 snellenella Wals., 488.  
 socors, 403.  
 solita, 45.  
 Sophronia, 33, 353.  
 sortifera, 557.  
 spanioctenis, 552.  
 sparactis, 440.  
 Sparganothis, 260, 370,  
     611.  
 sparsiciliella Clem.,  
     471.  
 Spatialistis, 116, 611.  
 spatulata, 405.  
 sphacterias, 383.  
 Sphaerelictis, 102.  
 sphaerobola, 120.  
 Sphalerostola, 365.  
 spiloglypta, 548.  
 Spilonota, 67.  
 spilopis, 356.  
 spilozeucta, 366.  
 spinigera, 5.

- spinulosa, 114.  
 Sporadartha, 320.  
 sporochlora, 523.  
 sporodesma, 464.  
 sporomochla, 491.  
 spumans, 605.  
 spumescens, 131.  
 spumosa, 260.  
 squalida, 272.  
 squamifera, 525.  
 Stachyostoma, 28.  
 Stagmatophora, 388,  
     546.  
 stagmatopis, 50.  
 Stagmaturgis, 25.  
 stagnicolor, 229.  
 staphidotis, 604.  
 Stasixena, 573.  
 Statherotis, 443.  
 Stathmopoda, 66, 96,  
     377, 541, 585.  
 stationaria, 402.  
 Stegasta, 18.  
 stelocosma, 145.  
 stenochorda, 456.  
 Stenolechia, 12, 484.  
 Stenoma, 178, 239,  
     365, 557.  
 stenomorpha, 49.  
 Stenoptilia, 301, 570.  
 Steremniodes, 37.  
 sterrhomitra, 178.  
 stibogramma, 90.  
 stibomorpha, 484.  
 Stichotactis, 562.  
 stigmatophora Wals.,  
     545.  
 stiphropa, 70.  
 Stiphrostola, 25.  
 stirpicola, 342.  
 stomobapta, 524.  
 Stomopteryx, 281, 352.  
 strepsibathra, 446.  
 stringens, 164.  
 strophopa, 46.  
 strophota, 252.  
 stygeropa, 187.  
 subagrestis, 573.  
 subcrocea, 60.  
 subdulcis, 195.  
 subflavescens, 568.  
 subita, 224.  
 sublimbata Zell., 205.  
 sublimicola, 626.  
 subridens, 64.  
 subrutila, 1.  
 subsidiaria, 107.  
 substrigata, 397.  
 substructa, 367.  
 summivola, 625.  
 superans, 291.  
 superciliaris, 577.  
 suprema, 301.  
 surda, 6.  
 suspensa, 19.  
 sustentata, 225.  
 symbolaspis, 334.  
 Symmoeca, 292, 356,  
     516.  
 symmochlopa, 519.  
 symmorias, 52.  
 symphanes, 264.  
 Symphorostola, 376.  
 sympiestis, 129.  
 symplacota, 606.  
 Synanthesdon, 267.  
 Syncathedra, 37.  
 synchrysa, 66.  
 syncolleta, 44.  
 Syncratomorpha, 509.  
 synerita, 281.  
 syncrotaula, 317.  
 syndyas, 286.  
 synercta, 173.  
 syngalactis, 389.  
 synglypta, 579.  
 syngonarcha, 290.  
 syntonopis, 48.  
 syringota, 287.  
 syrista, 293.  
 Syrmologa, 326.  
 syrphetopa, 278.  
 Syscalma, 619.  
 tabellifera, 379.  
 tabulatrix, 597.  
 Talaeporia, 75.  
 talaria, 77.  
 talarodes, 329.  
 Tanyarches, 99.  
 tanyphaea, 77.  
 tanysphena, 408.  
 tanysta, 189.  
 taospila, 243.  
 taphiopis, 485.  
 taphrocopa, 317.  
 taurina, 477.  
 technographa, 391.  
 telemacha, 356.  
 teleodelta, 295.  
 teleogramma, 128.  
 teleopa, 333.  
 teleosema, 165.  
 Telphusa, 16, 275, 344,  
     487.  
 temperatella Led., 61.  
 tepens, 2.  
 tephrinopa, 496.  
 tephroclysta, 553  
 tephroclystis, 616.  
 tephromorpha, 613.  
 tephroplintha, 30.  
 tepidus, 299.  
 terebrata, 87.  
 terenaula, 374.  
 teretracma, 353.  
 termanches, 500.  
 terpnodes, 15.  
 Terthreutis, 459.  
 tetrachorda, 91.  
 Tetracmanthes, 136.  
 tetradeta, 292.  
 tetralina, 627.  
 tetrameris, 277.  
 tetrapetra, 194.  
 tetraploca, 446.  
 tetrargyra, 473.  
 Tetraschalis, 92.  
 tetratypa, 410.  
 thalamectis, 521.  
 thalamias, 30.  
 Thamnocrana, 382.  
 Thamnoscelis, 466.  
 Theatrocopia, 577.  
 Thelyasceta, 27, 509.  
 Themeliotis, 549.  
 theodora, 265.  
 theriosema, 404.  
 thermoconis, 139.  
 thermogramma, 502  
 thermopetra, 602.  
 thermophaea, 3.  
 thermopsis, 50.  
 thermurga, 540.  
 thianthes, 375.  
 thinoclasta, 280.  
 thiostoma, 509.  
 Thiotricha, 24, 280,  
     499.  
 Thisizima, 428.  
 thoristes Busck, 192.  
 thrasydoxa, 297.  
 thrasymedes, 624.  
 thrylodes, 580.  
 thrypsandra, 5.  
 thyellopa, 298.  
 thymiata, 497.  
 thymophanes, 411.  
 thyrota, 516.  
 thrysogastra, 529.  
 Thrysostoma, 499.  
 Thysiarcha, 146.  
 thysiarcha, 469.  
 Timocratica, 176.  
 Timodora, 86.  
 Tinaegeria, 586.  
 Tinea, 71, 322, 332,  
     427, 547, 552.  
 Tineola, 71.

- Tinissa, 319, 424.  
 Tinthia, 268.  
 Tipulomima, 465.  
 Tiquadra, 73.  
 Tisis, 516.  
 titania, 405.  
 tolerans, 599.  
 tonans, 135.  
 Tonica, 105, 474.  
 tornogramma, 171.  
 tornograpta, 341.  
 tornophanes, 603.  
 torodoxa, 386.  
 torquata, 339.  
 torrens, 334.  
 torridula, 304.  
 Tortrix, 54, 114, 247,  
     369, 457, 608.  
 Tortyra, 302, 423.  
 toxocosma, 501.  
 toxogramma, 143.  
 Toxotacma, 504.  
 Trachybyrsis, 368.  
 trachycosma, 21.  
 trachydesma, 534.  
 Trachyedra, 497.  
 Trachypepla, 469.  
 trachyphloeus, 300.  
 tractaria, 172.  
 tranquilla, 383.  
 transformata, 579.  
 transfossa, 318.  
 transvecta, 377.  
 trapezantha, 321.  
 trapeziodes, 253.  
 trasias, 448.  
 triacmias, 304.  
 tribapta, 455.  
 tribolopa, 66.  
 tribolopis, 344.  
 tribrachia, 38.  
 tribrachynta, 388.  
 tricharacta, 119.  
 trichaspis, 484.  
 Trichembola, 271.  
 Trichoboscis, 526.  
 Trichocerota, 269.  
 trichocyma, 47.  
 trichodyta, 77.  
 trichonota, 235.  
 trichopla, 518.  
 Trichoptilus, 564.  
 trichostola, 22, 347.  
 Trichotaphe, 4, 287,  
     355, 512.  
 tricitra, 583.  
 Triclonella, 98, 467.  
 Tricyanaula, 508.  
 tridentifera, 328.  
 triexoda, 125.  
 trigama, 476.  
 triglenopa, 520.  
 trilobopa, 249.  
 trilychna, 394.  
 trimelaena, 598.  
 trinephela, 487.  
 triploca, 406.  
 tripudians, 65.  
 Trisophista, 117.  
 trithalama, 547.  
 Trithamnora, 324.  
 trithyris, 266.  
 tritogramma, 176.  
 trivinctella Zell., 412.  
 Trochilium, 266.  
 trychota, 510.  
 trymalopa, 211.  
 trypheropa, 81.  
 turbinaris, 435.  
 tylonota, 247.  
 tympanista, 479.  
 Tyrictaca, 372.  
 tyriochalca, 587.  
 tyrocrossa, 183.  
 tyroplaca, 134.  
 tyroxesta, 213.  
 ulmicola, 612.  
 ulosema, 559.  
 umbellifera, 254.  
 Untomia, 503.  
 uranauges, 266.  
 Urodus, 128.  
 Ussara, 307.  
 vacata, 170.  
 vacuefacta, 148.  
 vadosa, 153.  
 vasifera, 204.  
 vectigalis, 380.  
 veligera, 458.  
 velitans, 54.  
 venatrix, 612.  
 verna, 72.  
 versabilis, 251.  
 vestigata, 485.  
 vexillifera, 112.  
 vitrea, 158.  
 vitreola, 209.  
 vivida, 103, 313.  
 vociferans, 574.  
 volitans, 220.  
 vomerata, 88.  
 vorticosa, 551.  
 xanthocapna, 607.  
 xanthocausta, 544.  
 xanthocharis, 518.  
 xanthoclista, 50.  
 xanthocosma, 46.  
 xanthodora, 24.  
 xanthoglypta, 423.  
 xanthomorpha, 42.  
 xanthophracta, 624.  
 xanthoplaca Turn.,  
     132.  
 xanthoschema, 320.  
 xanthosoma Dogn.,  
     192.  
 xanthoteles, 38.  
 xanthotyla, 546.  
 xanthuris, 346.  
 Xenothictis, 609.  
 Xerantica, 553.  
 xestochalca, 248.  
 xuthosaris, 175.  
 xylinarcha, 589.  
 xylinopa, 203.  
 xylochroa, 275.  
 xylodelta, 444.  
 Xylodryas, 151.  
 xyloglypta, 22.  
 xylomorpha, 498.  
 xylophragma, 307.  
 Zacorisca, 111, 369.  
 Zalithia, 508.  
 zalomorpha, 550.  
 Zelleria, 414, 562, 591.  
 zelosaris, 12.  
 zephyritis, 218.  
 zesticopa, 514.  
 zetes, 569.  
 zonodeta, 288.  
 zonoreas, 79.  
 zonotorna, 131, 593.  
 zophocrossa, 530.  
 zymogramma, 260.

END OF VOL. III.

